


# 1b3o

Evolutionary trace report by **report\_maker**

May 2, 2010



4.3.1	<b>Alistat</b>	8
4.3.2	<b>CE</b>	8
4.3.3	<b>DSSP</b>	8
4.3.4	<b>HSSP</b>	8
4.3.5	<b>LaTex</b>	8
4.3.6	<b>Muscle</b>	8
4.3.7	<b>Pymol</b>	8
4.4	Note about ET Viewer	8
4.5	Citing this work	8
4.6	About report_maker	8
4.7	Attachments	8

## CONTENTS

### 1 Introduction

### 2 Chain 1b3oB

- 2.1 P12268 overview
- 2.2 Multiple sequence alignment for 1b3oB
- 2.3 Residue ranking in 1b3oB
- 2.4 Top ranking residues in 1b3oB and their position on the structure
  - 2.4.1 Clustering of residues at 25% coverage.
  - 2.4.2 Overlap with known functional surfaces at 25% coverage.
  - 2.4.3 Possible novel functional surfaces at 25% coverage.

### 3 Notes on using trace results

- 3.1 Coverage
- 3.2 Known substitutions
- 3.3 Surface
- 3.4 Number of contacts
- 3.5 Annotation
- 3.6 Mutation suggestions

### 4 Appendix

- 4.1 File formats
- 4.2 Color schemes used
- 4.3 Credits

## 1 INTRODUCTION

From the original Protein Data Bank entry (PDB id 1b3o):

**Title:** Ternary complex of human type-ii inosine monophosphate dehydrogenase with 6-cl-imp and selenazole adenine dinucleotide

**Compound:** Mol id: 1; molecule: protein (inosine monophosphate dehydrogenase 2); chain: a, b; synonym: impd, impdh; ec: 1.1.1.205; engineered: yes


**Organism, scientific name:** Homo Sapiens;

1b3o contains a single unique chain 1b3oB (414 residues long) and its homologue 1b3oA.


## 1 2 CHAIN 1B3OB

### 2.1 P12268 overview

- 1 From SwissProt, id P12268, 84% identical to 1b3oB:
- 1 **Description:** Inosine-5'-monophosphate dehydrogenase 2 (EC
- 1 1.1.1.205) (IMP dehydrogenase 2) (IMPDH-II) (IMPD 2).
- 1 **Organism, scientific name:** Homo sapiens (Human).
- 1 **Taxonomy:** Eukaryota; Metazoa; Chordata; Craniata;
- 2 Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates;
- 2 Catarrhini; Hominidae; Homo.
- 3 **Function:** Rate limiting enzyme in the de novo synthesis of guanine
- 6 nucleotides and therefore is involved in the regulation of cell growth.
- 6 It may also have a role in the development of malignancy and the
- 6 growth progression of some tumors.
- 7 **Catalytic activity:** Inosine 5'-phosphate + NAD(+) + H(2)O =
- 7 xanthosine 5'-phosphate + NADH.
- 7 **Pathway:** GMP biosynthesis from IMP; first step.
- 7 **Subunit:** Homotetramer.
- 7 **Tissue specificity:** IMP type I is the main species in normal leuko-
- 7 cytes and type II predominates over type I in the tumor.
- 7 **Ptm:** The N-terminus is blocked.
- 7 **Similarity:** Belongs to the IMPDH/GMPR family.
- 7 **Similarity:** Contains 2 CBS domains.
- 7 **About:** This Swiss-Prot entry is copyright. It is produced through a
- 7 collaboration between the Swiss Institute of Bioinformatics and the
- 8 EMBL outstation - the European Bioinformatics Institute. There are



**Fig. 1.** Residues 10-244 in 1b3oB colored by their relative importance. (See Appendix, Fig.11, for the coloring scheme.)



**Fig. 2.** Residues 245-499 in 1b3oB colored by their relative importance. (See Appendix, Fig.11, for the coloring scheme.)

no restrictions on its use as long as its content is in no way modified and this statement is not removed.

## 2.2 Multiple sequence alignment for 1b3oB

For the chain 1b3oB, the alignment 1b3oB.msf (attached) with 74 sequences was used. The alignment was assembled through combination of BLAST searching on the UniProt database and alignment using Muscle program. It can be found in the attachment to this report, under the name of 1b3oB.msf. Its statistics, from the *alistat* program are the following:

```

Format: MSF
Number of sequences: 74
Total number of residues: 29516
Smallest: 386
Largest: 414
Average length: 398.9
Alignment length: 414
Average identity: 49%
Most related pair: 99%
Most unrelated pair: 29%
Most distant seq: 50%

```

Furthermore, 8% of residues show as conserved in this alignment.

The alignment consists of 37% eukaryotic ( 10% vertebrata, 2% arthropoda, 13% fungi, 8% plantae), 58% prokaryotic, and 2% archaean sequences. (Descriptions of some sequences were not readily available.) The file containing the sequence descriptions can be found in the attachment, under the name 1b3oB.descr.


## 2.3 Residue ranking in 1b3oB

The 1b3oB sequence is shown in Figs. 1–2, with each residue colored according to its estimated importance. The full listing of residues in 1b3oB can be found in the file called 1b3oB.ranks\_sorted in the attachment.

## 2.4 Top ranking residues in 1b3oB and their position on the structure


In the following we consider residues ranking among top 25% of residues in the protein . Figure 3 shows residues in 1b3oB colored

by their importance: bright red and yellow indicate more conserved/important residues (see Appendix for the coloring scheme). A Pymol script for producing this figure can be found in the attachment.



**Fig. 3.** Residues in 1b3oB, colored by their relative importance. Clockwise: front, back, top and bottom views.

**2.4.1 Clustering of residues at 25% coverage.** Fig. 4 shows the top 25% of all residues, this time colored according to clusters they belong to. The clusters in Fig.4 are composed of the residues listed in Table 1.



**Fig. 4.** Residues in 1b3oB, colored according to the cluster they belong to: red, followed by blue and yellow are the largest clusters (see Appendix for the coloring scheme). Clockwise: front, back, top and bottom views. The corresponding Pymol script is attached.

Table 1.		
cluster color	size	member residues
red	77	31, 32, 33, 34, 36, 64, 67, 68, 70 71, 72, 73, 74, 75, 89, 92, 94, 100 107, 108, 244, 259, 263, 267, 269 274, 276, 277, 278, 280, 302, 303 305, 306, 313, 317, 321, 322, 324 325, 326, 328, 329, 330, 331, 332 333, 334, 336, 338, 340, 342, 343 346, 362, 363, 364, 365, 366, 371 372, 375, 376, 379, 380, 381, 384 385, 387, 391, 392, 395, 397, 398 490, 493, 497
blue	5	143, 154, 156, 179, 218
yellow	5	39, 44, 469, 470, 472
green	4	79, 83, 86, 87
purple	2	459, 463

**Table 1.** Clusters of top ranking residues in 1b3oB.

**2.4.2 Overlap with known functional surfaces at 25% coverage.** The name of the ligand is composed of the source PDB identifier and the heteroatom name used in that file.


**Unknown atom or ion binding site.** Table 2 lists the top 25% of residues at the interface with 1b3oUNX112 (unknown atom or ion). The following table (Table 3) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 2.					
res	type	subst's (%)	cvg	noc/bb	dist (Å)
332	I	T (70) I (29)	0.15	2/0	4.37

**Table 2.** The top 25% of residues in 1b3oB at the interface with unknown atom or ion. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 3.		
res	type	disruptive mutations
332	I	(R) (YH) (K) (E)

**Table 3.** List of disruptive mutations for the top 25% of residues in 1b3oB, that are at the interface with unknown atom or ion.



**Fig. 5.** Residues in 1b3oB, at the interface with unknown atom or ion, colored by their relative importance. The ligand (unknown atom or ion) is colored green. Atoms further than 30Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain 1b3oB.)

Figure 5 shows residues in 1b3oB colored by their importance, at the interface with 1b3oUNX112.

**CPR binding site.** Table 4 lists the top 25% of residues at the interface with 1b3oBCPR631 (cpr). The following table (Table 5) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
70	M	M(100)	0.09	15/0	3.84
303	N	N(100)	0.09	3/0	4.08
328	G	G(100)	0.09	24/24	2.93
329	S	S(100)	0.09	44/32	2.95
331	C	C(100)	0.09	22/9	1.77
364	D	D(100)	0.09	24/4	2.44
365	G	G(100)	0.09	17/17	3.43
366	G	G(100)	0.09	7/7	3.41
385	M	M(100)	0.09	3/1	3.93
387	G	G(100)	0.09	14/14	3.00
334	Q	R(62) Q(37)	0.11	16/6	3.69
322	R	K(59) R(40)	0.13	8/0	3.64
330	I	I(97) A(2)	0.14	8/8	4.32
332	I	T(70) I(29)	0.15	2/2	4.88
68	S	A(51) S(48)	0.22	7/0	3.32


**Table 4.** The top 25% of residues in 1b3oB at the interface with CPR.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/ bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand. )

res	type	disruptive mutations
70	M	(Y)(TH)(SCRG)(FWD)
303	N	(Y)(FTWH)(SEVCARG)(MD)
328	G	(KER)(FQMWH)(NYLPI)(SVA)
329	S	(KR)(FQMWH)(NYELPI)(D)
331	C	(KER)(FQMWH)(NYLPI)(SVA)
364	D	(R)(FWH)(KYVCAG)(TQM)
365	G	(KER)(FQMWH)(NYLPI)(SVA)
366	G	(KER)(FQMWH)(NYLPI)(SVA)
385	M	(Y)(TH)(SCRG)(FWD)
387	G	(KER)(FQMWH)(NYLPI)(SVA)
334	Q	(Y)(T)(FW)(SVCAHG)
322	R	(T)(YD)(SVCAG)(FELWPI)
330	I	(YR)(H)(TKE)(SQCDG)
332	I	(R)(YH)(K)(E)
68	S	(KR)(QH)(FYEMW)(N)

**Table 5.** List of disruptive mutations for the top 25% of residues in 1b3oB, that are at the interface with CPR.

Figure 6 shows residues in 1b3oB colored by their importance, at the interface with 1b3oBCPR631.

**Unknown atom or ion binding site.** Table 6 lists the top 25% of residues at the interface with 1b3oUNX113 (unknown atom or



**Fig. 6.** Residues in 1b3oB, at the interface with CPR, colored by their relative importance. The ligand (CPR) is colored green. Atoms further than 30Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain 1b3oB.)

ion). The following table (Table 7) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
94	N	N(98) F(1)	0.09	1/0	4.70


**Table 6.** The top 25% of residues in 1b3oB at the interface with unknown atom or ion.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/ bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand. )

res	type	disruptive mutations
94	N	(TY)(E)(SCHRG)(FVAWD)

**Table 7.** List of disruptive mutations for the top 25% of residues in 1b3oB, that are at the interface with unknown atom or ion.

Figure 7 shows residues in 1b3oB colored by their importance, at the interface with 1b3oUNX113.

**Interface with 1b3oB3.** Table 8 lists the top 25% of residues at the interface with 1b3oB3. The following table (Table 9) suggests possible disruptive replacements for these residues (see Section 3.6).



**Fig. 7.** Residues in 1b3oB, at the interface with unknown atom or ion, colored by their relative importance. The ligand (unknown atom or ion) is colored green. Atoms further than 30Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain 1b3oB.)

Table 8.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
34	D	D(100)	0.09	26/0	2.79
340	G	G(100)	0.09	38/38	2.80
31	T	T(98) A(1)	0.10	6/0	3.58
338	A	G(62) A(37)	0.11	45/34	2.79
306	T	T(97) S(2)	0.13	8/0	3.93
342	P	P(90) G(8) A(1)	0.13	15/2	3.34
336	V	V(95) M(1) I(2)	0.18	8/5	4.04
276	S	S(52) A(47)	0.20	2/0	4.73
280	S	S(93) N(5) Q(1)	0.21	32/22	2.92
305	V	A(60) V(39)	0.21	2/0	3.68
325	M	I(56) M(37) V(5)	0.21	8/0	4.26
277	Q	H(60) Q(33) N(5)	0.22	38/1	3.34
30	L	L(83) Y(10) F(2) M(1) V(1)	0.25	1/0	3.81


**Table 8.** The top 25% of residues in 1b3oB at the interface with 1b3oB3. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 9.		
res	type	disruptive mutations
34	D	(R)(FWH)(KYVCAG)(TQM)
340	G	(KER)(FQMWHD)(NYLPI)(SVA)
31	T	(KR)(QH)(FEMW)(N)
338	A	(KER)(Y)(QHD)(N)
306	T	(KR)(FQMWH)(NELPI)(D)
342	P	(R)(Y)(H)(KE)
336	V	(Y)(R)(H)(KE)
276	S	(KR)(QH)(FYEMW)(N)
280	S	(R)(FWH)(Y)(K)

*continued in next column*

res	type	disruptive mutations
305	V	(KYER) (QHD) (N) (FTMW)
325	M	(Y) (H) (T) (R)
277	Q	(Y) (T) (FW) (SVCAHG)
30	L	(R) (Y) (TK) (E)

**Table 9.** List of disruptive mutations for the top 25% of residues in 1b3oB, that are at the interface with 1b3oB3.



**Fig. 8.** Residues in 1b3oB, at the interface with 1b3oB3, colored by their relative importance. 1b3oB3 is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 1b3oB.)

Figure 8 shows residues in 1b3oB colored by their importance, at the interface with 1b3oB3.

**SAE binding site.** Table 10 lists the top 25% of residues at the interface with 1b3oSAE601 (sae). The following table (Table 11) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
274	D	D(100)	0.09	34/14	2.48
303	N	N(100)	0.09	13/0	2.88
324	G	G(100)	0.09	4/4	3.18
364	D	D(100)	0.09	3/0	3.74
334	Q	R(62) Q(37)	0.11	28/0	2.92
322	R	K(59)	0.13	4/0	4.12

*continued in next column*

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
276	S	R(40) S(52) A(47)	0.20	60/37	2.77
325	M	I(56) M(37) V(5)	0.21	3/2	4.47
277	Q	H(60) Q(33) N(5)	0.22	1/1	4.78

**Table 10.** The top 25% of residues in 1b3oB at the interface with SAE.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

res	type	disruptive mutations
274	D	(R) (FWH) (KYVCAG) (TQM)
303	N	(Y) (FTWH) (SEVCARG) (MD)
324	G	(KER) (FQMWH) (NYLPI) (SVA)
364	D	(R) (FWH) (KYVCAG) (TQM)
334	Q	(Y) (T) (FW) (SVCAHG)
322	R	(T) (YD) (SVCAG) (FELWPI)
276	S	(KR) (QH) (FYEMW) (N)
325	M	(Y) (H) (T) (R)
277	Q	(Y) (T) (FW) (SVCAHG)


**Table 11.** List of disruptive mutations for the top 25% of residues in 1b3oB, that are at the interface with SAE.

Figure 9 shows residues in 1b3oB colored by their importance, at the interface with 1b3oSAE601.


**2.4.3 Possible novel functional surfaces at 25% coverage.** One group of residues is conserved on the 1b3oB surface, away from (or substantially larger than) other functional sites and interfaces recognizable in PDB entry 1b3o. It is shown in Fig. 10. The residues belonging to this surface "patch" are listed in Table 12, while Table 13 suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	substitutions(%)	cvg
39	P	P(100)	0.09
472	G	G(100)	0.09
469	Q	G(59)Q(37)S(1) F(1)	0.15
470	D	L(14)Y(47)D(37)	0.22
44	F	V(56)F(37)L(1) I(4)	0.24

*continued in next column*



**Fig. 9.** Residues in 1b3oB, at the interface with SAE, colored by their relative importance. The ligand (SAE) is colored green. Atoms further than 30Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain 1b3oB.)



**Fig. 10.** A possible active surface on the chain 1b3oB.

Table 12. continued			
res	type	substitutions(%)	cvg
468	C	M(63)L(8)F(8) C(20)	0.25

**Table 12.** Residues forming surface "patch" in 1b3oB.

Table 13.		
res	type	disruptive mutations
39	P	(YR)(TH)(SKECG)(FQWD)
472	G	(KER)(FQMWH)(NYLPI)(SVA)
469	Q	(Y)(H)(FW)(T)
470	D	(R)(K)(H)(FW)
44	F	(KE)(TR)(QD)(SCG)
468	C	(R)(KE)(HD)(YQ)

**Table 13.** Disruptive mutations for the surface patch in 1b3oB.

### 3 NOTES ON USING TRACE RESULTS

#### 3.1 Coverage

Trace results are commonly expressed in terms of coverage: the residue is important if its "coverage" is small - that is if it belongs to some small top percentage of residues [100% is all of the residues in a chain], according to trace. The ET results are presented in the form of a table, usually limited to top 25% percent of residues (or to some nearby percentage), sorted by the strength of the presumed evolutionary pressure. (I.e., the smaller the coverage, the stronger the pressure on the residue.) Starting from the top of that list, mutating a couple of residues should affect the protein somehow, with the exact effects to be determined experimentally.

#### 3.2 Known substitutions

One of the table columns is "substitutions" - other amino acid types seen at the same position in the alignment. These amino acid types may be interchangeable at that position in the protein, so if one wants to affect the protein by a point mutation, they should be avoided. For example if the substitutions are "RVK" and the original protein has an R at that position, it is advisable to try anything, but RVK. Conversely, when looking for substitutions which will *not* affect the protein, one may try replacing, R with K, or (perhaps more surprisingly), with V. The percentage of times the substitution appears in the alignment is given in the immediately following bracket. No percentage is given in the cases when it is smaller than 1%. This is meant to be a rough guide - due to rounding errors these percentages often do not add up to 100%.

#### 3.3 Surface

To detect candidates for novel functional interfaces, first we look for residues that are solvent accessible (according to DSSP program) by at least 10Å<sup>2</sup>, which is roughly the area needed for one water molecule to come in the contact with the residue. Furthermore, we require that these residues form a "cluster" of residues which have neighbor within 5Å from any of their heavy atoms.

Note, however, that, if our picture of protein evolution is correct, the neighboring residues which *are not* surface accessible might be equally important in maintaining the interaction specificity - they should not be automatically dropped from consideration when choosing the set for mutagenesis. (Especially if they form a cluster with the surface residues.)

#### 3.4 Number of contacts

Another column worth noting is denoted "noc/bb"; it tells the number of contacts heavy atoms of the residue in question make across

the interface, as well as how many of them are realized through the backbone atoms (if all or most contacts are through the backbone, mutation presumably won't have strong impact). Two heavy atoms are considered to be "in contact" if their centers are closer than 5Å.

### 3.5 Annotation

If the residue annotation is available (either from the pdb file or from other sources), another column, with the header "annotation" appears. Annotations carried over from PDB are the following: site (indicating existence of related site record in PDB), S-S (disulfide bond forming residue), hb (hydrogen bond forming residue, jb (james bond forming residue), and sb (for salt bridge forming residue).

### 3.6 Mutation suggestions

Mutation suggestions are completely heuristic and based on complementarity with the substitutions found in the alignment. Note that they are meant to be **disruptive** to the interaction of the protein with its ligand. The attempt is made to complement the following properties: small [AVGSTC], medium [LPNQDEMIK], large [WFYHR], hydrophobic [LPVAMWFI], polar [GTCY]; positively [KHR], or negatively [DE] charged, aromatic [WFYH], long aliphatic chain [EK RQM], OH-group possession [SDETY], and NH2 group possession [NQRK]. The suggestions are listed according to how different they appear to be from the original amino acid, and they are grouped in round brackets if they appear equally disruptive. From left to right, each bracketed group of amino acid types resembles more strongly the original (i.e. is, presumably, less disruptive) These suggestions are tentative - they might prove disruptive to the fold rather than to the interaction. Many researcher will choose, however, the straightforward alanine mutations, especially in the beginning stages of their investigation.

## 4 APPENDIX

### 4.1 File formats

Files with extension "ranks.sorted" are the actual trace results. The fields in the table in this file:

- alignment# number of the position in the alignment
- residue# residue number in the PDB file
- type amino acid type
- rank rank of the position according to older version of ET
- variability has two subfields:
  1. number of different amino acids appearing in in this column of the alignment
  2. their type
- rho ET score - the smaller this value, the lesser variability of this position across the branches of the tree (and, presumably, the greater the importance for the protein)
- cvg coverage - percentage of the residues on the structure which have this rho or smaller
- gaps percentage of gaps in this column

### 4.2 Color schemes used

The following color scheme is used in figures with residues colored by cluster size: black is a single-residue cluster; clusters composed of more than one residue colored according to this hierarchy (ordered




Fig. 11. Coloring scheme used to color residues by their relative importance.

by descending size): red, blue, yellow, green, purple, azure, turquoise, brown, coral, magenta, LightSalmon, SkyBlue, violet, gold, bisque, LightSlateBlue, orchid, RosyBrown, MediumAquamarine, DarkOliveGreen, CornflowerBlue, grey55, burlywood, LimeGreen, tan, DarkOrange, DeepPink, maroon, BlanchedAlmond.

The colors used to distinguish the residues by the estimated evolutionary pressure they experience can be seen in Fig. 11.

### 4.3 Credits

**4.3.1 Alistat** *alistat* reads a multiple sequence alignment from the file and shows a number of simple statistics about it. These statistics include the format, the number of sequences, the total number of residues, the average and range of the sequence lengths, and the alignment length (e.g. including gap characters). Also shown are some percent identities. A percent pairwise alignment identity is defined as (idents / MIN(len1, len2)) where idents is the number of exact identities and len1, len2 are the unaligned lengths of the two sequences. The "average percent identity", "most related pair", and "most unrelated pair" of the alignment are the average, maximum, and minimum of all (N)(N-1)/2 pairs, respectively. The "most distant seq" is calculated by finding the maximum pairwise identity (best relative) for all N sequences, then finding the minimum of these N numbers (hence, the most outlying sequence). *alistat* is copyrighted by HHMI/Washington University School of Medicine, 1992-2001, and freely distributed under the GNU General Public License.

**4.3.2 CE** To map ligand binding sites from different source structures, *report\_maker* uses the CE program: <http://cl.sdsc.edu/>. Shindyalov IN, Bourne PE (1998) "Protein structure alignment by incremental combinatorial extension (CE) of the optimal path". *Protein Engineering* 11(9) 739-747.

**4.3.3 DSSP** In this work a residue is considered solvent accessible if the DSSP program finds it exposed to water by at least 10Å<sup>2</sup>, which is roughly the area needed for one water molecule to come in the contact with the residue. DSSP is copyrighted by W. Kabsch, C. Sander and MPI-MF, 1983, 1985, 1988, 1994 1995, CMBI version by Elmar.Krieger@cmbi.kun.nl November 18,2002,

<http://www.cmbi.kun.nl/gv/dssp/descrip.html>.



**4.3.4 HSSP** Whenever available, report\_maker uses HSSP alignment as a starting point for the analysis (sequences shorter than 75% of the query are taken out, however); R. Schneider, A. de Daruvar, and C. Sander. "The HSSP database of protein structure-sequence alignments." *Nucleic Acids Res.*, 25:226–230, 1997.

<http://swift.cmbi.kun.nl/swift/hssp/>

**4.3.5 LaTeX** The text for this report was processed using L<sup>A</sup>T<sub>E</sub>X; Leslie Lamport, "LaTeX: A Document Preparation System Addison-Wesley," Reading, Mass. (1986).

**4.3.6 Muscle** When making alignments "from scratch", report maker uses Muscle alignment program: Edgar, Robert C. (2004), "MUSCLE: multiple sequence alignment with high accuracy and high throughput." *Nucleic Acids Research* 32(5), 1792-97.

<http://www.drive5.com/muscle/>

**4.3.7 Pymol** The figures in this report were produced using Pymol. The scripts can be found in the attachment. Pymol is an open-source application copyrighted by DeLano Scientific LLC (2005). For more information about Pymol see <http://pymol.sourceforge.net/>. (Note for Windows users: the attached package needs to be unzipped for Pymol to read the scripts and launch the viewer.)

#### 4.4 Note about ET Viewer

Dan Morgan from the Lichtarge lab has developed a visualization tool specifically for viewing trace results. If you are interested, please visit:

<http://mammoth.bcm.tmc.edu/traceview/>

The viewer is self-unpacking and self-installing. Input files to be used with ETV (extension .etvx) can be found in the attachment to the main report.

#### 4.5 Citing this work

The method used to rank residues and make predictions in this report can be found in Mihalek, I., I. Reš, O. Lichtarge. (2004). "A Family of Evolution-Entropy Hybrid Methods for Ranking of Protein Residues

by Importance" *J. Mol. Bio.* **336**: 1265-82. For the original version of ET see O. Lichtarge, H. Bourne and F. Cohen (1996). "An Evolutionary Trace Method Defines Binding Surfaces Common to Protein Families" *J. Mol. Bio.* **257**: 342-358.

report\_maker itself is described in Mihalek I., I. Res and O. Lichtarge (2006). "Evolutionary Trace Report Maker: a new type of service for comparative analysis of proteins." *Bioinformatics* **22**:1656-7.

#### 4.6 About report\_maker

report\_maker was written in 2006 by Ivana Mihalek. The 1D ranking visualization program was written by Ivica Reš. report\_maker is copyrighted by Lichtarge Lab, Baylor College of Medicine, Houston.

#### 4.7 Attachments

The following files should accompany this report:

- 1b3oB.complex.pdb - coordinates of 1b3oB with all of its interacting partners
- 1b3oB.etvx - ET viewer input file for 1b3oB
- 1b3oB.cluster\_report.summary - Cluster report summary for 1b3oB
- 1b3oB.ranks - Ranks file in sequence order for 1b3oB
- 1b3oB.clusters - Cluster descriptions for 1b3oB
- 1b3oB.msf - the multiple sequence alignment used for the chain 1b3oB
- 1b3oB.descr - description of sequences used in 1b3oB msf
- 1b3oB.ranks.sorted - full listing of residues and their ranking for 1b3oB
- 1b3oB.1b3oUNX112.if.pml - Pymol script for Figure 5
- 1b3oB.cbvcg - used by other 1b3oB – related pymol scripts
- 1b3oB.1b3oBCPR631.if.pml - Pymol script for Figure 6
- 1b3oB.1b3oUNX113.if.pml - Pymol script for Figure 7
- 1b3oB.1b3oB3.if.pml - Pymol script for Figure 8
- 1b3oB.1b3oSAE601.if.pml - Pymol script for Figure 9