


# 1egc

Evolutionary trace report by **report\_maker**

March 9, 2010



## CONTENTS

### 1 Introduction

### 2 Chain 1egcA

- 2.1 Q5T4U4 overview
- 2.2 Multiple sequence alignment for 1egcA
- 2.3 Residue ranking in 1egcA
- 2.4 Top ranking residues in 1egcA and their position on the structure
  - 2.4.1 Clustering of residues at 25% coverage.
  - 2.4.2 Overlap with known functional surfaces at 25% coverage.

### 3 Notes on using trace results

- 3.1 Coverage
- 3.2 Known substitutions
- 3.3 Surface
- 3.4 Number of contacts
- 3.5 Annotation
- 3.6 Mutation suggestions

### 4 Appendix

- 4.1 File formats
- 4.2 Color schemes used
- 4.3 Credits
  - 4.3.1 **Alistat**
  - 4.3.2 **CE**

4.3.3	<b>DSSP</b>	12
4.3.4	<b>HSSP</b>	12
4.3.5	<b>LaTex</b>	12
4.3.6	<b>Muscle</b>	12
4.3.7	<b>Pymol</b>	12
4.4	Note about ET Viewer	12
4.5	Citing this work	12
4.6	About report_maker	13
4.7	Attachments	13

## 1 INTRODUCTION

From the original Protein Data Bank entry (PDB id 1egc):

**Title:** Structure of t255e, e376g mutant of human medium chain acyl-coa dehydrogenase complexed with octanoyl-coa

**Compound:** Mol id: 1; molecule: medium chain acyl-coa dehydrogenase; chain: a, b, c, d; ec: 1.3.99.3; engineered: yes; mutation: yes; other details: complexed with octanoyl-coa

**Organism, scientific name:** Homo Sapiens;

1 1egc contains a single unique chain 1egcA (387 residues long) and its homologues 1egcD, 1egcC, and 1egcB.

1

1

1

1

1

1

2

2

2

2

11

11

11

11

11

11

11

11

11

12

12

12

12

12

12

12

12

12

12

12

12

12

12

## 2 CHAIN 1EGCA

### 2.1 Q5T4U4 overview

From SwissProt, id Q5T4U4, 99% identical to 1egcA:


**Description:** Acyl-Coenzyme A dehydrogenase, C-4 to C-12 straight chain.

**Organism, scientific name:** Homo sapiens (Human).


**Taxonomy:** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Catarrhini; Hominidae; Homo.

### 2.2 Multiple sequence alignment for 1egcA

11 For the chain 1egcA, the alignment 1egcA.msf (attached) with 1164  
11 sequences was used. The alignment was downloaded from the HSSP  
12 database, and fragments shorter than 75% of the query as well as  
12 duplicate sequences were removed. It can be found in the attachment  
12 to this report, under the name of 1egcA.msf. Its statistics, from the  
12 *alistat* program are the following:



**Fig. 1.** Residues 10-202 in legcA colored by their relative importance. (See Appendix, Fig.13, for the coloring scheme.)



**Fig. 2.** Residues 203-396 in legcA colored by their relative importance. (See Appendix, Fig.13, for the coloring scheme.)

```

Format:                MSF
Number of sequences:   1164
Total number of residues: 438009
Smallest:              300
Largest:               387
Average length:        376.3
Alignment length:      387
Average identity:      40%
Most related pair:     99%
Most unrelated pair:   23%
Most distant seq:     38%

```

Furthermore, <1% of residues show as conserved in this alignment.

The alignment consists of 5% eukaryotic ( 3% vertebrata, <1% arthropoda, <1% fungi, <1% plantae), 14% prokaryotic, and 1% archaean sequences. (Descriptions of some sequences were not readily available.) The file containing the sequence descriptions can be found in the attachment, under the name legcA.descr.


### 2.3 Residue ranking in legcA

The legcA sequence is shown in Figs. 1–2, with each residue colored according to its estimated importance. The full listing of residues in legcA can be found in the file called legcA.ranks.sorted in the attachment.

### 2.4 Top ranking residues in legcA and their position on the structure

In the following we consider residues ranking among top 25% of residues in the protein . Figure 3 shows residues in legcA colored

by their importance: bright red and yellow indicate more conserved/important residues (see Appendix for the coloring scheme). A Pymol script for producing this figure can be found in the attachment.




**Fig. 3.** Residues in legcA, colored by their relative importance. Clockwise: front, back, top and bottom views.

2.4.1 Clustering of residues at 25% coverage. Fig. 4 shows the top 25% of all residues, this time colored according to clusters they belong to. The clusters in Fig.4 are composed of the residues listed in Table 1.

Table 1.		
cluster color	size	member residues
red	71	43,103,132,133,135,136,137 138,141,142,143,144,149,151 162,164,166,167,168,169,170 173,180,182,193,197,212,214 216,217,220,222,242,244,245 248,249,252,253,255,256,259 270,315,316,329,330,332,333 338,347,349,352,353,355,356 363,364,367,368,370,371,374 375,376,377,378,379,380,382 383
blue	4	85,86,265,309
yellow	3	283,284,285
green	2	71,72
purple	2	299,301
azure	2	112,116

*continued in next column*



**Fig. 4.** Residues in legcA, colored according to the cluster they belong to: red, followed by blue and yellow are the largest clusters (see Appendix for the coloring scheme). Clockwise: front, back, top and bottom views. The corresponding Pymol script is attached.

Table 1. continued		
cluster color	size	member residues
turquoise	2	277, 281

**Table 1.** Clusters of top ranking residues in legcA.

**2.4.2 Overlap with known functional surfaces at 25% coverage.** The name of the ligand is composed of the source PDB identifier and the heteroatom name used in that file.


**Octanoyl-coenzyme a binding site.** Table 2 lists the top 25% of residues at the interface with legcBCO8400 (octanoyl-coenzyme a). The following table (Table 3) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 2.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
284	F	F(97) YG M(1) SDN .	0.04	10/0	3.58

**Table 2.** The top 25% of residues in legcA at the interface with octanoyl-coenzyme a. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 3.		
res	type	disruptive mutations
284	F	(K) (E) (QR) (TD)

**Table 3.** List of disruptive mutations for the top 25% of residues in legcA, that are at the interface with octanoyl-coenzyme a.



**Fig. 5.** Residues in legcA, at the interface with octanoyl-coenzyme a, colored by their relative importance. The ligand (octanoyl-coenzyme a) is colored green. Atoms further than 30 Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain legcA.)

Figure 5 shows residues in legcA colored by their importance, at the interface with legcBCO8400.


**Interface with legcC.** Table 4 lists the top 25% of residues at the interface with legcC. The following table (Table 5) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 4.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
292	Q	Q(95) G P(1) Y E(1) TAS LH	0.07	55/10	2.90

**Table 4.** The top 25% of residues in legcA at the interface with legcC. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 5.		
res	type	disruptive mutations
292	Q	(Y) (H) (FW) (T)

**Table 5.** List of disruptive mutations for the top 25% of residues in legcA, that are at the interface with legcC.



**Fig. 6.** Residues in legcA, at the interface with legcC, colored by their relative importance. legcC is shown in backbone representation (See Appendix for the coloring scheme for the protein chain legcA.)

Figure 6 shows residues in legcA colored by their importance, at the interface with legcC.


**FAD binding site.** Table 6 lists the top 25% of residues at the interface with legcDFAD399 (fad). The following table (Table 7) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 6.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
292	Q	Q (95) G P (1) Y E (1) T A S L H	0.07	10/0	3.06

**Table 6.** The top 25% of residues in legcA at the interface with FAD. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 7.		
res	type	disruptive mutations
292	Q	(Y) (H) (FW) (T)

**Table 7.** List of disruptive mutations for the top 25% of residues in legcA, that are at the interface with FAD.



**Fig. 7.** Residues in legcA, at the interface with FAD, colored by their relative importance. The ligand (FAD) is colored green. Atoms further than 30Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain legcA.)

Figure 7 shows residues in legcA colored by their importance, at the interface with legcDFAD399.

**FAD binding site.** Table 8 lists the top 25% of residues at the interface with legcAFAD399 (fad). The following table (Table 9) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 8.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
141	G	G (100)	0.00	17/17	3.45
214	N	LK (94) N (3) TMR VF	0.04	2/0	4.17
376	G	E (78) A (17) G (3) TD. VS	0.04	1/1	4.70

*continued in next column*

Table 8. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
136	T	T(64) S(35)	0.07	39/9	2.71
142	S	S(93)	0.07	28/11	3.31
167	I	T(6) V(3) I(90) L(1)T A(1) C(1)SN	0.08	21/17	3.84
168	T	T(76) S(22)GH	0.09	32/16	2.69
222	T	V(2) T(91) S(3)EIN LAHQPMC D	0.09	11/0	3.65
375	Y	Y(49) L(13) G(27) F(5) V(1)W A(1).PI CE	0.09	31/3	3.61
374	I	I(94)F V(1) L(3)P.M T	0.10	13/9	3.53
380	Q	E(70) Q(18) D(6) N(3).AH ISR	0.10	32/6	2.62
378	T	T(93) A(3) S(3).V	0.11	14/1	2.72
135	V	L(66) M(20) V(5)TFY I(3)SAN C.	0.12	12/8	3.12
166	W	F(49) W(47) Y(3)L	0.13	62/34	2.77
103	L	V(9) I(10)A M(15) L(58) F(1)E C(1)DWN QGTYS	0.15	2/0	4.44
133	Y	F(45) I(2)	0.16	21/12	2.94

continued in next column

Table 8. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
217	Q	Y(26) M(1) L(19) W(1)GTV ASH M(20) I(27) L(11) W(17) Y(3) Q(15)S V(2)DNC HRFA	0.18	1/0	3.94
371	I	V(26) I(40) A(1) L(28) F(1)TGM Q.PSY	0.21	9/0	3.38
381	I	V(20) I(77) M(1)L.A S	0.25	8/1	4.59

**Table 8.** The top 25% of residues in legcA at the interface with FAD.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/ bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)


Table 9.		
res	type	disruptive mutations
141	G	(KER)(FQMWHD)(NYLPI)(SVA)
214	N	(Y)(T)(H)(FW)
376	G	(R)(K)(H)(FEW)
136	T	(KR)(FMWH)(Q)(LPI)
142	S	(KR)(FQMWH)(NELPI)(Y)
167	I	(R)(Y)(H)(K)
168	T	(K)(R)(QM)(FEW)
222	T	(R)(K)(H)(FW)
375	Y	(K)(QR)(E)(M)
374	I	(R)(Y)(H)(T)
380	Q	(Y)(H)(FTW)(CG)
378	T	(KR)(Q)(H)(M)
135	V	(R)(K)(E)(Y)
166	W	(K)(E)(Q)(TD)
103	L	(R)(Y)(H)(K)
133	Y	(K)(Q)(R)(E)
217	Q	(Y)(T)(H)(FW)

continued in next column



res	type	disruptive mutations
371	I	(R) (Y) (H) (K)
381	I	(YR) (H) (T) (K)

**Table 9.** List of disruptive mutations for the top 25% of residues in *legcA*, that are at the interface with FAD.



**Fig. 8.** Residues in *legcA*, at the interface with FAD, colored by their relative importance. The ligand (FAD) is colored green. Atoms further than 30 Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain *legcA*.)

Figure 8 shows residues in *legcA* colored by their importance, at the interface with *legcAFAD399*.

**Interface with *legcB*.** Table 10 lists the top 25% of residues at the interface with *legcB*. The following table (Table 11) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
216	G	G(99)V	0.01	11/11	3.00
281	R	R(99)QH	0.01	23/0	3.13
352	G	G(99)AS	0.02	11/11	3.64
143	D	D(95) E(2)NA	0.03	55/13	2.92
367	R	R(98)AN L.V	0.03	14/4	3.45
214	N	LK(94)	0.04	58/24	2.91

*continued in next column*

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
284	F	N(3)TMR VF F(97)YG M(1)SDN	0.04	46/7	3.56
349	Q	Q(92) E(2) R(3)HDN TLKG	0.05	52/4	2.76
353	G	G(96)P A(2)SID WYVM	0.05	19/19	3.31
142	S	S(93) T(6)	0.07	6/5	3.73
374	I	I(94)F V(1) L(3)P.M T	0.10	32/2	3.37
380	Q	E(70) Q(18) D(6) N(3).AH ISR	0.10	15/0	3.43
363	E	E(75)Q G(16) S(1) A(4)N.C H	0.11	21/3	3.43
378	T	T(93) A(3) S(3).V	0.11	2/0	4.41
370	K	R(39) K(55) P(2)HAE M.IWV	0.12	45/3	2.75
144	V	A(59) V(32) L(3)I P(2)NT S(1)M	0.13	1/1	4.89
166	W	F(49) W(47) Y(3)L	0.13	32/0	3.28
212	E	E(58) D(4) L(20) Y(5) K(1) H(1)VG I(2) F(2)QNP TAMR.	0.14	9/2	3.63

*continued in next column*

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
285	G	K(1) G(93) D(2)P N(2)EQR S.	0.14	1/1	4.94
356	F	Y(83) F(14)LI V(1)C	0.14	119/20	2.87
217	Q	M(20) I(27) L(11) W(17) Y(3) Q(15)S V(2)DNC HRFA	0.18	3/3	3.77
283	T	Q(64) A(20) S(3) T(10)EV ILRPD.	0.20	11/8	3.11
138	P	P(86) S(2) A(5) R(1)V T(1)HEQ F	0.21	4/4	3.28
371	I	V(26) I(40) A(1) L(28) F(1)TGM Q.PSY	0.21	29/7	3.72
379	S	S(54) N(31) T(6) A(1)K Q(1)G R(1)D.I EHLIC	0.23	4/2	4.02
381	I	V(20) I(77) M(1)L.A S	0.25	5/0	3.64

**Table 10.** The top 25% of residues in legcA at the interface with legcB. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/ bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

res	type	disruptive mutations
216	G	(KER)(QHD)(FYMW)(N)
281	R	(T)(D)(SYE)(VCAG)
352	G	(KR)(E)(QH)(FMW)
143	D	(R)(H)(FW)(Y)
367	R	(TYD)(E)(S)(CG)
214	N	(Y)(T)(H)(FW)
284	F	(K)(E)(QR)(TD)
349	Q	(Y)(FW)(H)(T)
353	G	(R)(K)(E)(H)
142	S	(KR)(FQMWH)(NELPI)(Y)
374	I	(R)(Y)(H)(T)
380	Q	(Y)(H)(FTW)(CG)
363	E	(FWH)(Y)(R)(VA)
378	T	(KR)(Q)(H)(M)
370	K	(Y)(T)(CG)(S)
144	V	(R)(Y)(K)(E)
166	W	(K)(E)(Q)(TD)
212	E	(H)(FW)(Y)(R)
285	G	(R)(FW)(H)(YE)
356	F	(K)(E)(QR)(D)
217	Q	(Y)(T)(H)(FW)
283	T	(R)(K)(H)(FW)
138	P	(Y)(R)(H)(T)
371	I	(R)(Y)(H)(K)
379	S	(R)(K)(FWH)(Y)
381	I	(YR)(H)(T)(K)

**Table 11.** List of disruptive mutations for the top 25% of residues in legcA, that are at the interface with legcB.


Figure 9 shows residues in legcA colored by their importance, at the interface with legcB.

**Octanoyl-coenzyme a binding site.** Table 12 lists the top 25% of residues at the interface with legcACO8400 (octanoyl-coenzyme a). The following table (Table 13) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)	antn
141	G	G(100)	0.00	7/7	3.63	
256	R	R(99)L	0.02	9/0	2.90	
377	G	G(99).	0.02	10/10	3.36	
143	D	D(95) E(2)NA	0.03	5/3	4.16	
376	G	E(78) A(17) G(3)TD. VS	0.04	22/22	3.04	
136	T	T(64) S(35).	0.07	2/0	4.00	
142	S	S(93) T(6)	0.07	22/21	3.15	

continued in next column





**Fig. 9.** Residues in legcA, at the interface with legcB, colored by their relative importance. legcB is shown in backbone representation (See Appendix for the coloring scheme for the protein chain legcA.)

Table 12. continued						
res	type	subst's (%)	cvg	noc/ bb	dist (Å)	antn
252	F	L(87) F(10)I V(1)SY	0.07	27/0	3.14	
168	T	T(76) S(22)GH	0.09	6/0	3.52	
375	Y	Y(49) L(13) G(27) F(5) V(1)W A(1).PI CE	0.09	32/6	3.57	
255	E	G(64) A(1)N T(8) E(20) S(3)DIV RQF	0.10	4/0	4.10	site
135	V	L(66) M(20) V(5)TFY I(3)SAN C.	0.12	2/0	4.83	
144	V	A(59) V(32) L(3)I P(2)NT	0.13	20/9	3.04	


*continued in next column*

Table 12. continued						
res	type	subst's (%)	cvg	noc/ bb	dist (Å)	antn
103	L	S(1)M V(9) I(10)A M(15) L(58) F(1)E C(1)DWN QGTYHS	0.15	15/0	3.39	
249	M	L(20) M(73) F(1) I(2)AVN TSRQ	0.15	42/7	3.03	
259	V	I(58) V(16) L(20) M(1)ACG TQNS	0.15	3/0	4.07	
133	Y	F(45) I(2) Y(26) M(1) L(19) W(1)GTV ASH	0.16	3/0	3.72	
245	F	L(7) F(62) Y(9) V(14)HI TA(2)DG MQSE	0.16	27/0	3.15	
253	D	D(68) E(13) N(10) A(2) S(1) T(1)P Q(1)LG	0.17	11/0	2.74	
381	I	V(20) I(77) M(1)L.A S	0.25	8/0	3.80	

**Table 12.** The top 25% of residues in legcA at the interface with octanoyl-coenzyme a.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand. )

Table 13.		
res	type	disruptive mutations
141	G	(KER) (FQMWH) (NYLPI) (SVA)
256	R	(T) (YD) (SECG) (VA)
377	G	(KER) (FQMWH) (NLPI) (Y)
143	D	(R) (H) (FW) (Y)
376	G	(R) (K) (H) (FEW)
136	T	(KR) (FMWH) (Q) (LPI)
142	S	(KR) (FQMWH) (NELPI) (Y)
252	F	(K) (E) (QR) (D)
168	T	(K) (R) (QM) (FEW)
375	Y	(K) (QR) (E) (M)
255	E	(H) (FW) (R) (Y)
135	V	(R) (K) (E) (Y)
144	V	(R) (Y) (K) (E)
103	L	(R) (Y) (H) (K)
249	M	(Y) (H) (T) (R)
259	V	(R) (Y) (KE) (H)
133	Y	(K) (Q) (R) (E)
245	F	(K) (E) (R) (Q)
253	D	(R) (H) (FW) (Y)
381	I	(YR) (H) (T) (K)

**Table 13.** List of disruptive mutations for the top 25% of residues in legcA, that are at the interface with octanoyl-coenzyme a.



**Fig. 10.** Residues in legcA, at the interface with octanoyl-coenzyme a, colored by their relative importance. The ligand (octanoyl-coenzyme a) is colored green. Atoms further than 30 Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain legcA.)

Figure 10 shows residues in legcA colored by their importance, at the interface with legcACO8400.

**FAD binding site.** Table 14 lists the top 25% of residues at the interface with legcBFAD399 (fad). The following table (Table 15) suggests possible disruptive replacements for these residues (see Section 3.6).


Table 14.					
res	type	subst's (%)	cvg	noc/bb	dist (Å)
281	R	R(99)QH	0.01	20/0	2.73
352	G	G(99)AS	0.02	15/15	4.02
284	F	F(97)YG M(1)SDN	0.04	17/1	2.96
349	Q	Q(92) E(2) R(3)HDN TLKG	0.05	16/10	2.91
353	G	G(96)P A(2)SID WYVM	0.05	14/14	3.03
277	Y	Y(95) H(1)L F(1)WRQ	0.06	3/0	4.21
356	F	Y(83) F(14)LI V(1)C	0.14	5/0	4.47
283	T	Q(64) A(20) S(3) T(10)EV ILRPD.	0.20	20/0	3.29

**Table 14.** The top 25% of residues in legcA at the interface with FAD.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 15.		
res	type	disruptive mutations
281	R	(T) (D) (SYE) (VCAG)
352	G	(KR) (E) (QH) (FMW)
284	F	(K) (E) (QR) (TD)
349	Q	(Y) (FW) (H) (T)
353	G	(R) (K) (E) (H)
277	Y	(K) (E) (Q) (M)
356	F	(K) (E) (QR) (D)
283	T	(R) (K) (H) (FW)

**Table 15.** List of disruptive mutations for the top 25% of residues in legcA, that are at the interface with FAD.

Figure 11 shows residues in legcA colored by their importance, at the interface with legcBFAD399.



**Fig. 11.** Residues in legcA, at the interface with FAD, colored by their relative importance. The ligand (FAD) is colored green. Atoms further than 30Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain legcA.)

**Interface with legcD.** Table 16 lists the top 25% of residues at the interface with legcD. The following table (Table 17) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 16.					
res	type	subst's (%)	cvg	noc/bb	dist (Å)
292	Q	Q(95)G P(1)Y E(1)TAS LH	0.07	3/0	3.54
299	A	A(96) S(1) G(1) V(1)TM	0.08	23/11	3.17
380	Q	E(70) Q(18) D(6) N(3).AH ISR	0.10	16/12	2.90
383	R	R(70) K(16) Q(5) L(4)NME T.H	0.20	45/6	3.03
381	I	V(20) I(77)	0.25	1/1	4.94

*continued in next column*

Table 16. continued					
res	type	subst's (%)	cvg	noc/bb	dist (Å)
386	V	M(1)L.A S I(83) M(2) T(1) V(9)S Q(1) A(1)L.	0.25	16/3	3.41

**Table 16.** The top 25% of residues in legcA at the interface with legcD. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 17.		
res	type	disruptive mutations
292	Q	(Y)(H)(FW)(T)
299	A	(R)(K)(YE)(H)
380	Q	(Y)(H)(FTW)(CG)
383	R	(T)(Y)(D)(CG)
381	I	(YR)(H)(T)(K)
386	V	(R)(Y)(K)(H)

**Table 17.** List of disruptive mutations for the top 25% of residues in legcA, that are at the interface with legcD.

Figure 12 shows residues in legcA colored by their importance, at the interface with legcD.


### 3 NOTES ON USING TRACE RESULTS

#### 3.1 Coverage

Trace results are commonly expressed in terms of coverage: the residue is important if its “coverage” is small - that is if it belongs to some small top percentage of residues [100% is all of the residues in a chain], according to trace. The ET results are presented in the form of a table, usually limited to top 25% percent of residues (or to some nearby percentage), sorted by the strength of the presumed evolutionary pressure. (I.e., the smaller the coverage, the stronger the pressure on the residue.) Starting from the top of that list, mutating a couple of residues should affect the protein somehow, with the exact effects to be determined experimentally.

#### 3.2 Known substitutions

One of the table columns is “substitutions” - other amino acid types seen at the same position in the alignment. These amino acid types may be interchangeable at that position in the protein, so if one wants to affect the protein by a point mutation, they should be avoided. For example if the substitutions are “RVK” and the original protein has an R at that position, it is advisable to try anything, but RVK. Conversely, when looking for substitutions which will *not* affect the protein, one may try replacing, R with K, or (perhaps more surprisingly), with



**Fig. 12.** Residues in legcA, at the interface with legcD, colored by their relative importance. legcD is shown in backbone representation (See Appendix for the coloring scheme for the protein chain legcA.)

V. The percentage of times the substitution appears in the alignment is given in the immediately following bracket. No percentage is given in the cases when it is smaller than 1%. This is meant to be a rough guide - due to rounding errors these percentages often do not add up to 100%.

### 3.3 Surface

To detect candidates for novel functional interfaces, first we look for residues that are solvent accessible (according to DSSP program) by at least  $10\text{\AA}^2$ , which is roughly the area needed for one water molecule to come in the contact with the residue. Furthermore, we require that these residues form a “cluster” of residues which have neighbor within  $5\text{\AA}$  from any of their heavy atoms.

Note, however, that, if our picture of protein evolution is correct, the neighboring residues which *are not* surface accessible might be equally important in maintaining the interaction specificity - they should not be automatically dropped from consideration when choosing the set for mutagenesis. (Especially if they form a cluster with the surface residues.)

### 3.4 Number of contacts

Another column worth noting is denoted “noc/bb”; it tells the number of contacts heavy atoms of the residue in question make across the interface, as well as how many of them are realized through the backbone atoms (if all or most contacts are through the backbone, mutation presumably won’t have strong impact). Two heavy atoms are considered to be “in contact” if their centers are closer than  $5\text{\AA}$ .

### 3.5 Annotation

If the residue annotation is available (either from the pdb file or from other sources), another column, with the header “annotation” appears. Annotations carried over from PDB are the following: site

(indicating existence of related site record in PDB ), S-S (disulfide bond forming residue), hb (hydrogen bond forming residue, jb (james bond forming residue), and sb (for salt bridge forming residue).

### 3.6 Mutation suggestions

Mutation suggestions are completely heuristic and based on complementarity with the substitutions found in the alignment. Note that they are meant to be **disruptive** to the interaction of the protein with its ligand. The attempt is made to complement the following properties: small [AVGSTC], medium [LPNQDEMIK], large [WIFYHR], hydrophobic [LPVAMWFI], polar [GTCY]; positively [KHR], or negatively [DE] charged, aromatic [WIFYH], long aliphatic chain [EKRQM], OH-group possession [SDETY], and NH2 group possession [NQRK]. The suggestions are listed according to how different they appear to be from the original amino acid, and they are grouped in round brackets if they appear equally disruptive. From left to right, each bracketed group of amino acid types resembles more strongly the original (i.e. is, presumably, less disruptive) These suggestions are tentative - they might prove disruptive to the fold rather than to the interaction. Many researcher will choose, however, the straightforward alanine mutations, especially in the beginning stages of their investigation.

## 4 APPENDIX

### 4.1 File formats

Files with extension “ranks\_sorted” are the actual trace results. The fields in the table in this file:

- alignment# number of the position in the alignment
- residue# residue number in the PDB file
- type amino acid type
- rank rank of the position according to older version of ET
- variability has two subfields:
  1. number of different amino acids appearing in in this column of the alignment
  2. their type
- rho ET score - the smaller this value, the lesser variability of this position across the branches of the tree (and, presumably, the greater the importance for the protein)
- cvg coverage - percentage of the residues on the structure which have this rho or smaller
- gaps percentage of gaps in this column

### 4.2 Color schemes used

The following color scheme is used in figures with residues colored by cluster size: black is a single-residue cluster; clusters composed of more than one residue colored according to this hierarchy (ordered by descending size): red, blue, yellow, green, purple, azure, turquoise, brown, coral, magenta, LightSalmon, SkyBlue, violet, gold, bisque, LightSlateBlue, orchid, RosyBrown, MediumAquamarine, DarkOliveGreen, CornflowerBlue, grey55, burlywood, LimeGreen, tan, DarkOrange, DeepPink, maroon, BlanchedAlmond.

The colors used to distinguish the residues by the estimated evolutionary pressure they experience can be seen in Fig. 13.






Fig. 13. Coloring scheme used to color residues by their relative importance.

## 4.3 Credits

**4.3.1 Alistat** *alistat* reads a multiple sequence alignment from the file and shows a number of simple statistics about it. These statistics include the format, the number of sequences, the total number of residues, the average and range of the sequence lengths, and the alignment length (e.g. including gap characters). Also shown are some percent identities. A percent pairwise alignment identity is defined as  $(\text{idents} / \text{MIN}(\text{len1}, \text{len2}))$  where *idents* is the number of exact identities and *len1*, *len2* are the unaligned lengths of the two sequences. The "average percent identity", "most related pair", and "most unrelated pair" of the alignment are the average, maximum, and minimum of all  $(N)(N-1)/2$  pairs, respectively. The "most distant seq" is calculated by finding the maximum pairwise identity (best relative) for all *N* sequences, then finding the minimum of these *N* numbers (hence, the most outlying sequence). *alistat* is copyrighted by HHMI/Washington University School of Medicine, 1992-2001, and freely distributed under the GNU General Public License.

**4.3.2 CE** To map ligand binding sites from different source structures, *report\_maker* uses the CE program: <http://cl.sdsc.edu/>. Shindyalov IN, Bourne PE (1998) "Protein structure alignment by incremental combinatorial extension (CE) of the optimal path". *Protein Engineering* 11(9) 739-747.

**4.3.3 DSSP** In this work a residue is considered solvent accessible if the DSSP program finds it exposed to water by at least  $10\text{\AA}^2$ , which is roughly the area needed for one water molecule to come in the contact with the residue. DSSP is copyrighted by W. Kabsch, C. Sander and MPI-MF, 1983, 1985, 1988, 1994 1995, CMBI version by Elmar.Krieger@cmbi.kun.nl November 18,2002,

<http://www.cmbi.kun.nl/gv/dssp/descrip.html>.

**4.3.4 HSSP** Whenever available, *report\_maker* uses HSSP alignment as a starting point for the analysis (sequences shorter than 75% of the query are taken out, however); R. Schneider, A. de Daruvar, and C. Sander. "The HSSP database of protein structure-sequence alignments." *Nucleic Acids Res.*, 25:226-230, 1997.

<http://swift.cmbi.kun.nl/swift/hssp/>

**4.3.5 LaTeX** The text for this report was processed using  $\text{\LaTeX}$ ; Leslie Lamport, "LaTeX: A Document Preparation System Addison-Wesley," Reading, Mass. (1986).

**4.3.6 Muscle** When making alignments "from scratch", *report\_maker* uses Muscle alignment program: Edgar, Robert C. (2004), "MUSCLE: multiple sequence alignment with high accuracy and high throughput." *Nucleic Acids Research* 32(5), 1792-97.

<http://www.drive5.com/muscle/>

**4.3.7 Pymol** The figures in this report were produced using Pymol. The scripts can be found in the attachment. Pymol is an open-source application copyrighted by DeLano Scientific LLC (2005). For more information about Pymol see <http://pymol.sourceforge.net/>. (Note for Windows users: the attached package needs to be unzipped for Pymol to read the scripts and launch the viewer.)

## 4.4 Note about ET Viewer

Dan Morgan from the Lichtarge lab has developed a visualization tool specifically for viewing trace results. If you are interested, please visit:

<http://mammoth.bcm.tmc.edu/traceview/>

The viewer is self-unpacking and self-installing. Input files to be used with ETV (extension .etvx) can be found in the attachment to the main report.

## 4.5 Citing this work

The method used to rank residues and make predictions in this report can be found in Mihalek, I., I. Reš, O. Lichtarge. (2004). "A Family of Evolution-Entropy Hybrid Methods for Ranking of Protein Residues by Importance" *J. Mol. Bio.* **336**: 1265-82. For the original version of ET see O. Lichtarge, H.Bourne and F. Cohen (1996). "An Evolutionary Trace Method Defines Binding Surfaces Common to Protein Families" *J. Mol. Bio.* **257**: 342-358.

*report\_maker* itself is described in Mihalek I., I. Res and O. Lichtarge (2006). "Evolutionary Trace Report Maker: a new type of service for comparative analysis of proteins." *Bioinformatics* **22**:1656-7.

## 4.6 About report\_maker

*report\_maker* was written in 2006 by Ivana Mihalek. The 1D ranking visualization program was written by Ivica Reš. *report\_maker* is copyrighted by Lichtarge Lab, Baylor College of Medicine, Houston.

## 4.7 Attachments

The following files should accompany this report:

- *legcA.complex.pdb* - coordinates of *legcA* with all of its interacting partners
- *legcA.etvx* - ET viewer input file for *legcA*
- *legcA.cluster\_report.summary* - Cluster report summary for *legcA*
- *legcA.ranks* - Ranks file in sequence order for *legcA*
- *legcA.clusters* - Cluster descriptions for *legcA*
- *legcA.msf* - the multiple sequence alignment used for the chain *legcA*

- legcA.descr - description of sequences used in legcA msf
- legcA.ranks\_sorted - full listing of residues and their ranking for legcA
- legcA.legcBCO8400.if.pml - Pymol script for Figure 5
- legcA.cbcvg - used by other legcA – related pymol scripts
- legcA.legcC.if.pml - Pymol script for Figure 6
- legcA.legcDFAD399.if.pml - Pymol script for Figure 7
- legcA.legcAFAD399.if.pml - Pymol script for Figure 8
- legcA.legcB.if.pml - Pymol script for Figure 9
- legcA.legcACO8400.if.pml - Pymol script for Figure 10
- legcA.legcBFAD399.if.pml - Pymol script for Figure 11
- legcA.legcD.if.pml - Pymol script for Figure 12