


# 1g3j

Evolutionary trace report by **report\_maker**

September 9, 2008



## CONTENTS

### 1 Introduction

### 2 Chain 1g3jA

- 2.1 Q5R2I4 overview
- 2.2 Multiple sequence alignment for 1g3jA
- 2.3 Residue ranking in 1g3jA
- 2.4 Top ranking residues in 1g3jA and their position on the structure
  - 2.4.1 Clustering of residues at 26% coverage.
  - 2.4.2 Overlap with known functional surfaces at 26% coverage.
  - 2.4.3 Possible novel functional surfaces at 26% coverage.

### 3 Chain 1g3jB

- 3.1 P70063 overview
- 3.2 Multiple sequence alignment for 1g3jB
- 3.3 Residue ranking in 1g3jB
- 3.4 Top ranking residues in 1g3jB and their position on the structure
  - 3.4.1 Clustering of residues at 83% coverage.

### 4 Notes on using trace results

- 4.1 Coverage
- 4.2 Known substitutions
- 4.3 Surface

4.4	Number of contacts	7
4.5	Annotation	7
4.6	Mutation suggestions	7

## 5 Appendix **8**

5.1	File formats	8
5.2	Color schemes used	8
5.3	Credits	8
5.3.1	<b>Alistat</b>	8
5.3.2	<b>CE</b>	8
5.3.3	<b>DSSP</b>	8
5.3.4	<b>HSSP</b>	8
5.3.5	<b>LaTeX</b>	8
5.3.6	<b>Muscle</b>	8
5.3.7	<b>Pymol</b>	8
5.4	Note about ET Viewer	9
5.5	Citing this work	9
5.6	About report_maker	9
5.7	Attachments	9

## 1 INTRODUCTION

From the original Protein Data Bank entry (PDB id 1g3j):

**Title:** Crystal structure of the xtcf3-cbd/beta-catenin armadillo repeat complex

**Compound:** Mol id: 1; molecule: beta-catenin armadillo repeat region; chain: a, c; engineered: yes; mol id: 2; molecule: tcf3-cbd (catenin binding domain); chain: b, d; engineered: yes

**Organism, scientific name:** *Xenopus laevis*;

1g3j contains unique chains 1g3jA (522 residues) and 1g3jB (41 residues) 1g3jC is a homologue of chain 1g3jA. 1g3jD is a homologue of chain 1g3jB.

## 2 CHAIN 1G3JA

### 2.1 Q5R2I4 overview

From SwissProt, id Q5R2I4, 95% identical to 1g3jA:


**Description:** Beta-catenin homologue.

**Organism, scientific name:** *Trionyx sinensis* (Chinese softshell turtle) (*Pelodiscus sinensis*).


**Taxonomy:** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Testudines; Cryptodira; Trionychoidea; Trionychidae; Pelodiscus.

### 2.2 Multiple sequence alignment for 1g3jA

For the chain 1g3jA, the alignment 1g3jA.msf (attached) with 19 sequences was used. The alignment was assembled through combination of BLAST searching on the UniProt database and alignment using Muscle program. It can be found in the attachment to this



**Fig. 1.** Residues 134-394 in 1g3jA colored by their relative importance. (See Appendix, Fig.12, for the coloring scheme.)



**Fig. 2.** Residues 395-664 in 1g3jA colored by their relative importance. (See Appendix, Fig.12, for the coloring scheme.)

report, under the name of 1g3jA.msf. Its statistics, from the *alifold* program are the following:

```

Format:                MSF
Number of sequences:   19
Total number of residues: 9830
Smallest:              491
Largest:               522
Average length:        517.4
Alignment length:      522
Average identity:       57%
Most related pair:     99%
Most unrelated pair:   21%
Most distant seq:      34%

```

Furthermore, 6% of residues show as conserved in this alignment.

The alignment consists of 94% eukaryotic ( 36% vertebrata, 21% arthropoda) sequences. (Descriptions of some sequences were not readily available.) The file containing the sequence descriptions can be found in the attachment, under the name 1g3jA.descr.


### 2.3 Residue ranking in 1g3jA

The 1g3jA sequence is shown in Figs. 1–2, with each residue colored according to its estimated importance. The full listing of residues in 1g3jA can be found in the file called 1g3jA.ranks.sorted in the attachment.

### 2.4 Top ranking residues in 1g3jA and their position on the structure

In the following we consider residues ranking among top 26% of residues in the protein (the closest this analysis allows us to get to 25%). Figure 3 shows residues in 1g3jA colored by their importance: bright red and yellow indicate more conserved/important residues (see Appendix for the coloring scheme). A Pymol script for producing this figure can be found in the attachment.

**2.4.1 Clustering of residues at 26% coverage.** Fig. 4 shows the top 26% of all residues, this time colored according to clusters they belong to. The clusters in Fig.4 are composed of the residues listed




**Fig. 3.** Residues in 1g3jA, colored by their relative importance. Clockwise: front, back, top and bottom views.

in Table 1.

Table 1.		
cluster color	size	member residues
red	109	192, 195, 197, 222, 227, 229, 231, 232, 234, 235, 236, 237, 238, 240, 244, 250, 251, 254, 255, 257, 258, 259, 261, 262, 263, 286, 290, 292, 294, 295, 299, 301, 304, 312, 316, 320, 321, 324, 325, 333, 334, 335, 336, 339, 342, 345, 347, 348, 349

*continued in next column*



**Fig. 4.** Residues in 1g3jA, colored according to the cluster they belong to: red, followed by blue and yellow are the largest clusters (see Appendix for the coloring scheme). Clockwise: front, back, top and bottom views. The corresponding Pymol script is attached.

Table 1. continued		
cluster color	size	member residues
blue	15	350, 353, 354, 355, 358, 361, 365
		366, 370, 374, 376, 377, 382, 386
		387, 388, 389, 390, 393, 405, 419
		420, 422, 424, 425, 426, 427, 428
		429, 430, 431, 434, 435, 438, 443
		447, 462, 463, 466, 467, 468, 469
		470, 472, 474, 475, 481, 482, 505
		509, 512, 515, 516, 518, 519, 563
		564, 566, 571, 572
		577, 578, 581, 597, 599, 600, 602
yellow	3	603, 612, 618, 621, 640, 654, 655
		659
		167, 168, 171

**Table 1.** Clusters of top ranking residues in 1g3jA.

2.4.2 *Overlap with known functional surfaces at 26% coverage.* The name of the ligand is composed of the source PDB identifier and the heteroatom name used in that file.

**Interface with 1g3jB.** Table 2 lists the top 26% of residues at the interface with 1g3jB. The following table (Table 3) suggests possible disruptive replacements for these residues (see Section 4.6).

Table 2.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
292	K	K(100)	0.07	34/7	3.26
312	K	K(100)	0.07	9/0	2.72
376	R	R(100)	0.07	4/0	3.89
386	R	R(100)	0.07	45/8	3.47
387	N	N(100)	0.07	48/11	2.82
389	S	S(100)	0.07	2/0	4.33
390	D	D(100)	0.07	43/9	3.07
426	N	N(100)	0.07	41/3	2.84
435	K	K(100)	0.07	8/0	2.69
462	E	E(100)	0.07	5/1	4.46
469	R	R(100)	0.07	32/2	3.44
470	H	H(100)	0.07	47/9	3.29
250	S	S(94)	0.11	9/3	3.54
		K(5)			
254	Y	Y(94)	0.11	20/2	2.95
		C(5)			
299	D	D(94)	0.11	7/0	3.59
		N(5)			
425	S	S(94)	0.11	4/0	3.78
		C(5)			
430	N	N(94)	0.11	24/7	3.04
		G(5)			
466	C	C(94)	0.11	3/0	4.09
		A(5)			
519	L	L(94)	0.11	17/0	3.38
		M(5)			
261	N	N(89)	0.19	11/3	2.74
		S(10)			
335	K	K(89)	0.19	15/0	3.07
		N(10)			
349	V	V(89)	0.19	30/3	3.52
		.(10)			
354	K	K(89)	0.19	2/0	4.74
		A(10)			
422	G	G(89)	0.19	9/9	3.75
		Q(10)			
428	T	T(89)	0.19	3/3	4.04
		V(10)			
463	P	P(89)	0.19	4/2	4.10
		S(10)			
512	G	G(89)	0.19	5/5	3.60
		K(10)			
515	R	R(89)	0.19	57/0	3.34
		S(10)			
516	N	N(89)	0.19	22/1	2.88
		Q(10)			
571	E	E(89)	0.19	29/0	3.16
		H(10)			
612	R	R(89)	0.19	34/0	3.15

*continued in next column*

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
654	Y	S(10) Y(89) L(10)	0.19	10/0	2.79
393	T	T(84) S(15)	0.20	1/0	3.95
333	Y	Y(89) H(10)	0.22	4/0	3.67
345	K	K(89) R(10)	0.22	15/2	3.36


**Table 2.** The top 26% of residues in 1g3jA at the interface with 1g3jB. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

res	type	disruptive mutations
292	K	(Y)(FTW)(SVCAG)(HD)
312	K	(Y)(FTW)(SVCAG)(HD)
376	R	(TD)(SYEVCLAPIG)(FMW)(N)
386	R	(TD)(SYEVCLAPIG)(FMW)(N)
387	N	(Y)(FTWH)(SEVCARG)(MD)
389	S	(KR)(FQMWH)(NYELPI)(D)
390	D	(R)(FWH)(KYVCAG)(TQM)
426	N	(Y)(FTWH)(SEVCARG)(MD)
435	K	(Y)(FTW)(SVCAG)(HD)
462	E	(FWH)(YVCARG)(T)(SNKLPI)
469	R	(TD)(SYEVCLAPIG)(FMW)(N)
470	H	(E)(TQMD)(SNKVCLAPIG)(YR)
250	S	(FW)(YHR)(KM)(EQLPI)
254	Y	(K)(QM)(ER)(NLPI)
299	D	(R)(FWH)(Y)(VCAG)
425	S	(KR)(FQMWH)(E)(NYLPI)
430	N	(Y)(FWH)(ER)(T)
466	C	(KER)(QHD)(FYMW)(N)
519	L	(Y)(R)(TH)(SCG)
261	N	(Y)(FWH)(R)(TE)
335	K	(Y)(FTW)(SVCAG)(H)
349	V	(KYER)(QHD)(N)(FTMW)
354	K	(Y)(FTW)(SCHDG)(EVA)
422	G	(FEWHR)(KYD)(M)(QLPI)
428	T	(KR)(QH)(FEMW)(N)
463	P	(R)(Y)(H)(K)
512	G	(FEW)(YHDR)(KM)(QLPI)
515	R	(D)(TYELPI)(FVMCAWG)(S)
516	N	(Y)(FTWH)(SVCAG)(ER)
571	E	(FVCAWG)(TYHR)(SNKLPI)(QM)

*continued in next column*

res	type	disruptive mutations
612	R	(D)(TYELPI)(FVMCAWG)(S)
654	Y	(K)(QR)(EM)(NVA)
393	T	(KR)(FQMWH)(NELPI)(D)
333	Y	(K)(QM)(E)(NVLAPI)
345	K	(Y)(T)(FW)(SVCAG)

**Table 3.** List of disruptive mutations for the top 26% of residues in 1g3jA, that are at the interface with 1g3jB.



**Fig. 5.** Residues in 1g3jA, at the interface with 1g3jB, colored by their relative importance. 1g3jB is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 1g3jA.)

Figure 5 shows residues in 1g3jA colored by their importance, at the interface with 1g3jB.

**2.4.3 Possible novel functional surfaces at 26% coverage.** One group of residues is conserved on the 1g3jA surface, away from (or substantially larger than) other functional sites and interfaces recognizable in PDB entry 1g3j. It is shown in Fig. 6. The right panel shows (in blue) the rest of the larger cluster this surface belongs to. The residues belonging to this surface "patch" are listed in Table 4, while Table 5 suggests possible disruptive replacements for these residues (see Section 4.6).

res	type	substitutions(%)	cvg
192	P	P(94)F(5)	0.11
234	S	S(84)A(15)	0.20
235	G	G(84)D(15)	0.20

*continued in next column*



**Fig. 6.** A possible active surface on the chain 1g3jA. The larger cluster it belongs to is shown in blue.


Table 4. <i>continued</i>			
res	type	substitutions(%)	cvg
237	I	I(89)L(10)	0.22
238	P	P(89)A(10)	0.22

**Table 4.** Residues forming surface "patch" in 1g3jA.

Table 5.		
res	type	disruptive mutations
192	P	(R)(TY)(KE)(SCHG)
234	S	(KR)(QH)(FYEMW)(N)
235	G	(R)(K)(FWH)(EQM)
237	I	(YR)(TH)(SKECG)(FQWD)
238	P	(YR)(H)(TKE)(SQCDG)

**Table 5.** Disruptive mutations for the surface patch in 1g3jA.

Another group of surface residues is shown in Fig.7. The right panel shows (in blue) the rest of the larger cluster this surface belongs to.



**Fig. 7.** Another possible active surface on the chain 1g3jA. The larger cluster it belongs to is shown in blue.

The residues belonging to this surface "patch" are listed in Table 6, while Table 7 suggests possible disruptive replacements for these residues (see Section 4.6).

Table 6.			
res	type	substitutions(%)	cvg
292	K	K(100)	0.07
376	R	R(100)	0.07
335	K	K(89)N(10)	0.19
333	Y	Y(89)H(10)	0.22
334	E	E(89)R(10)	0.22
374	S	S(89)K(10)	0.22

**Table 6.** Residues forming surface "patch" in 1g3jA.

Table 7.		
res	type	disruptive mutations
292	K	(Y)(FTW)(SVCAG)(HD)
376	R	(TD)(SYEVCLAPIG)(FMW)(N)
335	K	(Y)(FTW)(SVCAG)(H)
333	Y	(K)(QM)(E)(NVLAPI)
334	E	(FW)(YVCAHG)(T)(SLPIR)
374	S	(FW)(YHR)(KM)(EQLPI)

**Table 7.** Disruptive mutations for the surface patch in 1g3jA.

Another group of surface residues is shown in Fig.8. The right panel shows (in blue) the rest of the larger cluster this surface belongs to.



**Fig. 8.** Another possible active surface on the chain 1g3jA. The larger cluster it belongs to is shown in blue.

The residues belonging to this surface "patch" are listed in Table 8, while Table 9 suggests possible disruptive replacements for these residues (see Section 4.6).

Table 8.			
res	type	substitutions(%)	cvg
312	K	K(100)	0.07
355	P	P(100)	0.07
361	G	G(100)	0.07
386	R	R(100)	0.07
387	N	N(100)	0.07
390	D	D(100)	0.07
426	N	N(100)	0.07

*continued in next column*

res	type	substitutions(%)	cvg
431	N	N(100)	0.07
434	N	N(100)	0.07
435	K	K(100)	0.07
462	E	E(100)	0.07
469	R	R(100)	0.07
470	H	H(100)	0.07
563	G	G(100)	0.07
564	V	V(100)	0.07
321	P	P(94)S(5)	0.11
425	S	S(94)C(5)	0.11
430	N	N(94)G(5)	0.11
466	C	C(94)A(5)	0.11
475	H	H(94)N(5)	0.11
482	Q	Q(94)V(5)	0.11
519	L	L(94)M(5)	0.11
659	L	L(94)M(5)	0.11
316	L	L(89)V(10)	0.19
349	V	V(89).(10)	0.19
350	C	C(89).(10)	0.19
354	K	K(89)A(10)	0.19
365	A	A(89)I(10)	0.19
419	C	C(89)Y(10)	0.19
422	G	G(89)Q(10)	0.19
463	P	P(89)S(10)	0.19
505	P	P(89)H(10)	0.19
509	A	A(89)L(10)	0.19
512	G	G(89)K(10)	0.19
515	R	R(89)S(10)	0.19
516	N	N(89)Q(10)	0.19
571	E	E(89)H(10)	0.19
612	R	R(89)S(10)	0.19
654	Y	Y(89)L(10)	0.19
393	T	T(84)S(15)	0.20
572	G	G(84)L(15)	0.20
345	K	K(89)R(10)	0.22
358	V	V(89)I(10)	0.22

Table 8. Residues forming surface "patch" in 1g3jA.

res	type	disruptive mutations
469	R	(TD)(SYEVCLAPIG)(FMW)(N)
470	H	(E)(TQMD)(SNKVCLAPIG)(YR)
563	G	(KER)(FQMWH)(NYLPI)(SVA)
564	V	(KYER)(QHD)(N)(FTMW)
321	P	(R)(Y)(H)(K)
425	S	(KR)(FQMH)(E)(NYLPI)
430	N	(Y)(FWH)(ER)(T)
466	C	(KER)(QHD)(FYMW)(N)
475	H	(E)(T)(MD)(SVQCAG)
482	Q	(Y)(H)(FTW)(SCDG)
519	L	(Y)(R)(TH)(SCG)
659	L	(Y)(R)(TH)(SCG)
316	L	(YR)(H)(TKE)(SQCDG)
349	V	(KYER)(QHD)(N)(FTMW)
350	C	(KER)(FQMWH)(NLPI)(Y)
354	K	(Y)(FTW)(SCHDG)(EVA)
365	A	(YR)(KE)(H)(QD)
419	C	(K)(ER)(QM)(D)
422	G	(FEWHR)(KYD)(M)(QLPI)
463	P	(R)(Y)(H)(K)
505	P	(TYR)(E)(SKCG)(QHD)
509	A	(YR)(KE)(H)(QD)
512	G	(FEW)(YHDR)(KM)(QLPI)
515	R	(D)(TYELPI)(FVMCAWG)(S)
516	N	(Y)(FTWH)(SVCAG)(ER)
571	E	(FVCAWG)(TYHR)(SNKLPI)(QM)
612	R	(D)(TYELPI)(FVMCAWG)(S)
654	Y	(K)(QR)(EM)(NVA)
393	T	(KR)(FQMH)(NELPI)(D)
572	G	(R)(KE)(H)(FYQWD)
345	K	(Y)(T)(FW)(SVCAG)
358	V	(YR)(KE)(H)(QD)

Table 9. Disruptive mutations for the surface patch in 1g3jA.

res	type	disruptive mutations
312	K	(Y)(FTW)(SVCAG)(HD)
355	P	(YR)(TH)(SKECG)(FQWD)
361	G	(KER)(FQMWH)(NYLPI)(SVA)
386	R	(TD)(SYEVCLAPIG)(FMW)(N)
387	N	(Y)(FTWH)(SEVCARG)(MD)
390	D	(R)(FWH)(KYVCAG)(TQM)
426	N	(Y)(FTWH)(SEVCARG)(MD)
431	N	(Y)(FTWH)(SEVCARG)(MD)
434	N	(Y)(FTWH)(SEVCARG)(MD)
435	K	(Y)(FTW)(SVCAG)(HD)
462	E	(FWH)(YVCARG)(T)(SNKLPI)

*continued in next column*

### 3 CHAIN 1G3JB

#### 3.1 P70063 overview

From SwissProt, id P70063, 82% identical to 1g3jB:

**Description:** Transcription factor XTFC-3c.

**Organism, scientific name:** *Xenopus laevis* (African clawed frog).

**Taxonomy:** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Amphibia; Batrachia; Anura; Mesobatrachia; Pipoidae; Pipidae; Xenopodinae; *Xenopus*; *Xenopus*.

#### 3.2 Multiple sequence alignment for 1g3jB

For the chain 1g3jB, the alignment 1g3jB.msf (attached) with 5 sequences was used. The alignment was assembled through combination of BLAST searching on the UniProt database and alignment using Muscle program. It can be found in the attachment to this report, under the name of 1g3jB.msf. Its statistics, from the *alistat* program are the following:



**Fig. 9.** Residues 2-52 in 1g3jB colored by their relative importance. (See Appendix, Fig.12, for the coloring scheme.)

```

Format:                MSF
Number of sequences:  5
Total number of residues: 205
Smallest:             41
Largest:              41
Average length:       41.0
Alignment length:     41
Average identity:     91%
Most related pair:    98%
Most unrelated pair:  88%
Most distant seq:     90%

```

Furthermore, 82% of residues show as conserved in this alignment.

The alignment consists of 80% eukaryotic ( 80% vertebrata) sequences. (Descriptions of some sequences were not readily available.) The file containing the sequence descriptions can be found in the attachment, under the name 1g3jB.descr.

### 3.3 Residue ranking in 1g3jB

The 1g3jB sequence is shown in Fig. 9, with each residue colored according to its estimated importance. The full listing of residues in 1g3jB can be found in the file called 1g3jB.ranks\_sorted in the attachment.


### 3.4 Top ranking residues in 1g3jB and their position on the structure

In the following we consider residues ranking among top 83% of residues in the protein (the closest this analysis allows us to get to 25%). Figure 10 shows residues in 1g3jB colored by their importance: bright red and yellow indicate more conserved/important residues (see Appendix for the coloring scheme). A Pymol script for producing this figure can be found in the attachment.


*3.4.1 Clustering of residues at 83% coverage.* Fig. 11 shows the top 83% of all residues, this time colored according to clusters they belong to. The clusters in Fig.11 are composed of the residues listed in Table 10.

Table 10.		
cluster color	size	member residues
red	22	2, 3, 4, 7, 8, 9, 10, 12, 13, 14, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 29
blue	12	40, 41, 42, 43, 45, 46, 47, 48, 49, 50, 51, 52

**Table 10.** Clusters of top ranking residues in 1g3jB.



**Fig. 10.** Residues in 1g3jB, colored by their relative importance. Clockwise: front, back, top and bottom views.



**Fig. 11.** Residues in 1g3jB, colored according to the cluster they belong to: red, followed by blue and yellow are the largest clusters (see Appendix for the coloring scheme). Clockwise: front, back, top and bottom views. The corresponding Pymol script is attached.

## 4 NOTES ON USING TRACE RESULTS

### 4.1 Coverage

Trace results are commonly expressed in terms of coverage: the residue is important if its “coverage” is small - that is if it belongs to some small top percentage of residues [100% is all of the residues

in a chain], according to trace. The ET results are presented in the form of a table, usually limited to top 25% percent of residues (or to some nearby percentage), sorted by the strength of the presumed evolutionary pressure. (I.e., the smaller the coverage, the stronger the pressure on the residue.) Starting from the top of that list, mutating a couple of residues should affect the protein somehow, with the exact effects to be determined experimentally.

## 4.2 Known substitutions

One of the table columns is “substitutions” - other amino acid types seen at the same position in the alignment. These amino acid types may be interchangeable at that position in the protein, so if one wants to affect the protein by a point mutation, they should be avoided. For example if the substitutions are “RVK” and the original protein has an R at that position, it is advisable to try anything, but RVK. Conversely, when looking for substitutions which will *not* affect the protein, one may try replacing R with K, or (perhaps more surprisingly), with V. The percentage of times the substitution appears in the alignment is given in the immediately following bracket. No percentage is given in the cases when it is smaller than 1%. This is meant to be a rough guide - due to rounding errors these percentages often do not add up to 100%.

## 4.3 Surface

To detect candidates for novel functional interfaces, first we look for residues that are solvent accessible (according to DSSP program) by at least  $10\text{\AA}^2$ , which is roughly the area needed for one water molecule to come in the contact with the residue. Furthermore, we require that these residues form a “cluster” of residues which have neighbor within  $5\text{\AA}$  from any of their heavy atoms.

Note, however, that, if our picture of protein evolution is correct, the neighboring residues which *are not* surface accessible might be equally important in maintaining the interaction specificity - they should not be automatically dropped from consideration when choosing the set for mutagenesis. (Especially if they form a cluster with the surface residues.)

## 4.4 Number of contacts

Another column worth noting is denoted “noc/bb”; it tells the number of contacts heavy atoms of the residue in question make across the interface, as well as how many of them are realized through the backbone atoms (if all or most contacts are through the backbone, mutation presumably won’t have strong impact). Two heavy atoms are considered to be “in contact” if their centers are closer than  $5\text{\AA}$ .

## 4.5 Annotation

If the residue annotation is available (either from the pdb file or from other sources), another column, with the header “annotation” appears. Annotations carried over from PDB are the following: site (indicating existence of related site record in PDB), S-S (disulfide bond forming residue), hb (hydrogen bond forming residue), jb (james bond forming residue), and sb (for salt bridge forming residue).

## 4.6 Mutation suggestions

Mutation suggestions are completely heuristic and based on complementarity with the substitutions found in the alignment. Note that they are meant to be **disruptive** to the interaction of the protein with its ligand. The attempt is made to complement the following properties: small [AVGSTC], medium [LPNQDEMILK], large




Fig. 12. Coloring scheme used to color residues by their relative importance.

[WFYHR], hydrophobic [LPVAMWFI], polar [GTCY]; positively [KHR], or negatively [DE] charged, aromatic [WFYH], long aliphatic chain [EK RQM], OH-group possession [SDETY], and NH2 group possession [NQRK]. The suggestions are listed according to how different they appear to be from the original amino acid, and they are grouped in round brackets if they appear equally disruptive. From left to right, each bracketed group of amino acid types resembles more strongly the original (i.e. is, presumably, less disruptive) These suggestions are tentative - they might prove disruptive to the fold rather than to the interaction. Many researcher will choose, however, the straightforward alanine mutations, especially in the beginning stages of their investigation.

## 5 APPENDIX

### 5.1 File formats

Files with extension “ranks\_sorted” are the actual trace results. The fields in the table in this file:

- alignment# number of the position in the alignment
- residue# residue number in the PDB file
- type amino acid type
- rank rank of the position according to older version of ET
- variability has two subfields:
  1. number of different amino acids appearing in in this column of the alignment
  2. their type
- rho ET score - the smaller this value, the lesser variability of this position across the branches of the tree (and, presumably, the greater the importance for the protein)
- cvg coverage - percentage of the residues on the structure which have this rho or smaller
- gaps percentage of gaps in this column

### 5.2 Color schemes used

The following color scheme is used in figures with residues colored by cluster size: black is a single-residue cluster; clusters composed of



more than one residue colored according to this hierarchy (ordered by descending size): red, blue, yellow, green, purple, azure, turquoise, brown, coral, magenta, LightSalmon, SkyBlue, violet, gold, bisque, LightSlateBlue, orchid, RosyBrown, MediumAquaMarine, DarkOliveGreen, CornflowerBlue, grey55, burlywood, LimeGreen, tan, DarkOrange, DeepPink, maroon, BlanchedAlmond.

The colors used to distinguish the residues by the estimated evolutionary pressure they experience can be seen in Fig. 12.

### 5.3 Credits

**5.3.1 Alistat** *alistat* reads a multiple sequence alignment from the file and shows a number of simple statistics about it. These statistics include the format, the number of sequences, the total number of residues, the average and range of the sequence lengths, and the alignment length (e.g. including gap characters). Also shown are some percent identities. A percent pairwise alignment identity is defined as  $(\text{idents} / \text{MIN}(\text{len1}, \text{len2}))$  where idents is the number of exact identities and len1, len2 are the unaligned lengths of the two sequences. The "average percent identity", "most related pair", and "most unrelated pair" of the alignment are the average, maximum, and minimum of all  $(N)(N-1)/2$  pairs, respectively. The "most distant seq" is calculated by finding the maximum pairwise identity (best relative) for all N sequences, then finding the minimum of these N numbers (hence, the most outlying sequence). *alistat* is copyrighted by HHMI/Washington University School of Medicine, 1992-2001, and freely distributed under the GNU General Public License.

**5.3.2 CE** To map ligand binding sites from different source structures, *report\_maker* uses the CE program: <http://cl.sdsc.edu/>. Shindyalov IN, Bourne PE (1998) "Protein structure alignment by incremental combinatorial extension (CE) of the optimal path". *Protein Engineering* 11(9) 739-747.

**5.3.3 DSSP** In this work a residue is considered solvent accessible if the DSSP program finds it exposed to water by at least  $10\text{\AA}^2$ , which is roughly the area needed for one water molecule to come in the contact with the residue. DSSP is copyrighted by W. Kabsch, C. Sander and MPI-MF, 1983, 1985, 1988, 1994 1995, CMBI version by Elmar.Krieger@cmbi.kun.nl November 18,2002,

<http://www.cmbi.kun.nl/gv/dssp/descrip.html>.

**5.3.4 HSSP** Whenever available, *report\_maker* uses HSSP alignment as a starting point for the analysis (sequences shorter than 75% of the query are taken out, however); R. Schneider, A. de Daruvar, and C. Sander. "The HSSP database of protein structure-sequence alignments." *Nucleic Acids Res.*, 25:226-230, 1997.

<http://swift.cmbi.kun.nl/swift/hssp/>

**5.3.5 LaTeX** The text for this report was processed using  $\text{\LaTeX}$ ; Leslie Lamport, "LaTeX: A Document Preparation System Addison-Wesley," Reading, Mass. (1986).

**5.3.6 Muscle** When making alignments "from scratch", *report\_maker* uses Muscle alignment program: Edgar, Robert C. (2004), "MUSCLE: multiple sequence alignment with high accuracy and high throughput." *Nucleic Acids Research* 32(5), 1792-97.

<http://www.drive5.com/muscle/>

**5.3.7 Pymol** The figures in this report were produced using Pymol. The scripts can be found in the attachment. Pymol

is an open-source application copyrighted by DeLano Scientific LLC (2005). For more information about Pymol see <http://pymol.sourceforge.net/>. (Note for Windows users: the attached package needs to be unzipped for Pymol to read the scripts and launch the viewer.)

### 5.4 Note about ET Viewer

Dan Morgan from the Lichtarge lab has developed a visualization tool specifically for viewing trace results. If you are interested, please visit:

<http://mammoth.bcm.tmc.edu/traceview/>

The viewer is self-unpacking and self-installing. Input files to be used with ETV (extension .etvx) can be found in the attachment to the main report.

### 5.5 Citing this work

The method used to rank residues and make predictions in this report can be found in Mihalek, I., I. Reš, O. Lichtarge. (2004). "A Family of Evolution-Entropy Hybrid Methods for Ranking of Protein Residues by Importance" *J. Mol. Bio.* **336**: 1265-82. For the original version of ET see O. Lichtarge, H.Bourne and F. Cohen (1996). "An Evolutionary Trace Method Defines Binding Surfaces Common to Protein Families" *J. Mol. Bio.* **257**: 342-358.

*report\_maker* itself is described in Mihalek I., I. Res and O. Lichtarge (2006). "Evolutionary Trace Report Maker: a new type of service for comparative analysis of proteins." *Bioinformatics* **22**:1656-7.

### 5.6 About report\_maker

*report\_maker* was written in 2006 by Ivana Mihalek. The 1D ranking visualization program was written by Ivica Reš. *report\_maker* is copyrighted by Lichtarge Lab, Baylor College of Medicine, Houston.

### 5.7 Attachments

The following files should accompany this report:

- 1g3jA.complex.pdb - coordinates of 1g3jA with all of its interacting partners
- 1g3jA.etvx - ET viewer input file for 1g3jA
- 1g3jA.cluster\_report.summary - Cluster report summary for 1g3jA
- 1g3jA.ranks - Ranks file in sequence order for 1g3jA
- 1g3jA.clusters - Cluster descriptions for 1g3jA
- 1g3jA.msf - the multiple sequence alignment used for the chain 1g3jA
- 1g3jA.descr - description of sequences used in 1g3jA msf
- 1g3jA.ranks\_sorted - full listing of residues and their ranking for 1g3jA
- 1g3jA.1g3jB.if.pml - Pymol script for Figure 5
- 1g3jA.cbv - used by other 1g3jA - related pymol scripts
- 1g3jB.complex.pdb - coordinates of 1g3jB with all of its interacting partners
- 1g3jB.etvx - ET viewer input file for 1g3jB

- 1g3jB.cluster\_report.summary - Cluster report summary for 1g3jB
- 1g3jB.ranks - Ranks file in sequence order for 1g3jB
- 1g3jB.clusters - Cluster descriptions for 1g3jB
- 1g3jB.msf - the multiple sequence alignment used for the chain 1g3jB
- 1g3jB.descr - description of sequences used in 1g3jB msf
- 1g3jB.ranks\_sorted - full listing of residues and their ranking for 1g3jB