


1oih

Evolutionary trace report by **report_maker**

July 13, 2010



CONTENTS

1 Introduction

2 Chain 1oihA

- 2.1 Q9WWU5 overview
- 2.2 Multiple sequence alignment for 1oihA
- 2.3 Residue ranking in 1oihA
- 2.4 Top ranking residues in 1oihA and their position on the structure
 - 2.4.1 Clustering of residues at 25% coverage.
 - 2.4.2 Overlap with known functional surfaces at 25% coverage.

3 Notes on using trace results

- 3.1 Coverage
- 3.2 Known substitutions
- 3.3 Surface
- 3.4 Number of contacts
- 3.5 Annotation
- 3.6 Mutation suggestions

4 Appendix

- 4.1 File formats
- 4.2 Color schemes used
- 4.3 Credits
 - 4.3.1 **Alistat**
 - 4.3.2 **CE**

4.3.3	DSSP	7
4.3.4	HSSP	7
4.3.5	LaTex	7
4.3.6	Muscle	7
4.3.7	Pymol	7
4.4	Note about ET Viewer	7
4.5	Citing this work	7
4.6	About report_maker	7
4.7	Attachments	7

1 INTRODUCTION

From the original Protein Data Bank entry (PDB id 1oih):

Title: Crystal structure of the alkylsulfatase atsk, a non-heme fe(ii) alphaketoglutarate dependent dioxygenase

Compound: Mol id: 1; molecule: putative alkylsulfatase atsk; chain: a, b, c, d

Organism, scientific name: Pseudomonas Putida;

1 1oih contains a single unique chain 1oihA (254 residues long) and its homologues 1oihB, 1oihD, and 1oihC.

1

1

1

1

2 CHAIN 1OIH A

2.1 Q9WWU5 overview

2 From SwissProt, id Q9WWU5, 88% identical to 1oihA:

Description: Putative alkylsulfatase.

Organism, scientific name: Pseudomonas putida.

Taxonomy: Bacteria; Proteobacteria; Gammaproteobacteria; Pseudomonadales; Pseudomonadaceae; Pseudomonas.

5

5

6

6

6

2.2 Multiple sequence alignment for 1oihA

6 For the chain 1oihA, the alignment 1oihA.msf (attached) with 466 sequences was used. The alignment was downloaded from the HSSP database, and fragments shorter than 75% of the query as well as duplicate sequences were removed. It can be found in the attachment to this report, under the name of 1oihA.msf. Its statistics, from the *alistat* program are the following:

7




Fig. 1. Residues 13-148 in 1oihA colored by their relative importance. (See Appendix, Fig.9, for the coloring scheme.)




Fig. 2. Residues 149-300 in 1oihA colored by their relative importance. (See Appendix, Fig.9, for the coloring scheme.)

```

Format: MSF
Number of sequences: 466
Total number of residues: 82396
Smallest: 96
Largest: 254
Average length: 176.8
Alignment length: 254
Average identity: 37%
Most related pair: 99%
Most unrelated pair: 0%
Most distant seq: 33%

```

Furthermore, <1% of residues show as conserved in this alignment.

The alignment consists of 9% eukaryotic (9% fungi), and 12% prokaryotic sequences. (Descriptions of some sequences were not readily available.) The file containing the sequence descriptions can be found in the attachment, under the name 1oihA.descr.

2.3 Residue ranking in 1oihA

The 1oihA sequence is shown in Figs. 1–2, with each residue colored according to its estimated importance. The full listing of residues in 1oihA can be found in the file called 1oihA.ranks.sorted in the attachment.

2.4 Top ranking residues in 1oihA and their position on the structure

In the following we consider residues ranking among top 25% of residues in the protein . Figure 3 shows residues in 1oihA colored by their importance: bright red and yellow indicate more conserved/important residues (see Appendix for the coloring scheme). A Pymol script for producing this figure can be found in the attachment.

2.4.1 Clustering of residues at 25% coverage. Fig. 4 shows the top 25% of all residues, this time colored according to clusters they belong to. The clusters in Fig.4 are composed of the residues listed




Fig. 3. Residues in 1oihA, colored by their relative importance. Clockwise: front, back, top and bottom views.




Fig. 4. Residues in 1oihA, colored according to the cluster they belong to: red, followed by blue and yellow are the largest clusters (see Appendix for the coloring scheme). Clockwise: front, back, top and bottom views. The corresponding Pymol script is attached.

in Table 1.

cluster color	size	member residues
red	63	107,108,110,111,112,113,117 118,121,123,129,132,133,134 135,137,139,143,144,147,158 160,162,196,197,198,199,200 202,203,205,206,207,208,210 212,213,215,219,221,226,227 230,231,234,246,247,249,253 255,257,258,259,260,264,266 269,275,277,279,281,284,287

Table 1. Clusters of top ranking residues in 1oihA.

2.4.2 *Overlap with known functional surfaces at 25% coverage.* The name of the ligand is composed of the source PDB identifier and the heteroatom name used in that file.

Sodium ion binding site. By analogy with 1oihC – 1oihCNA1300 interface. Table 2 lists the top 25% of residues at the interface with 1oihCNA1300 (sodium ion). The following table (Table 3) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
264	H	H(95) . (2)RDL ST	0.02	3/0	3.23
279	R	R(95)K . (3)HAS Q	0.03	1/0	4.07
108	H	H(77) . (22)	0.06	5/0	2.65
110	D	D(76) . (22)G	0.07	3/0	2.88

Table 2. The top 25% of residues in 1oihA at the interface with sodium ion.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

res	type	disruptive mutations
264	H	(E)(M)(Q)(TD)
279	R	(T)(D)(Y)(E)
108	H	(E)(TQMD)(SNVCLAPIG)(K)
110	D	(R)(FWH)(K)(VA)

Table 3. List of disruptive mutations for the top 25% of residues in 1oihA, that are at the interface with sodium ion.

Figure 5 shows residues in 1oihA colored by their importance, at the interface with 1oihCNA1300.




Fig. 5. Residues in 1oihA, at the interface with sodium ion, colored by their relative importance. The ligand (sodium ion) is colored green. Atoms further than 30 Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain 1oihA.)

Sodium ion binding site. Table 4 lists the top 25% of residues at the interface with 1oihANA1301 (sodium ion). The following table (Table 5) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
257	W	W(94) F(3) . (1)Y	0.01	1/0	4.90
264	H	H(95) . (2)RDL ST	0.02	5/0	2.24
279	R	R(95)K . (3)HAS Q	0.03	1/0	4.65
108	H	H(77) . (22)	0.06	5/0	2.33
110	D	D(76) . (22)G	0.07	5/1	2.25

Table 4. The top 25% of residues in 1oihA at the interface with sodium ion.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 5.		
res	type	disruptive mutations
257	W	(K) (E) (Q) (D)
264	H	(E) (M) (Q) (TD)
279	R	(T) (D) (Y) (E)
108	H	(E) (TQMD) (SNVCLAPIG) (K)
110	D	(R) (FWH) (K) (VA)

Table 5. List of disruptive mutations for the top 25% of residues in 1oihA, that are at the interface with sodium ion.




Fig. 6. Residues in 1oihA, at the interface with sodium ion, colored by their relative importance. The ligand (sodium ion) is colored green. Atoms further than 30Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain 1oihA.)

Figure 6 shows residues in 1oihA colored by their importance, at the interface with 1oihANA1301.

Interface with 1oihB. Table 6 lists the top 25% of residues at the interface with 1oihB. The following table (Table 7) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 6.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
221	G	D(7) N(9) G(68) E(9) Q(2)	0.09	4/4	3.51

continued in next column

Table 6. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
226	D	.(1) S(1)P E(75) D(9) Q(2) G(4) A(2) . (1)RVT KMHFNL	0.15	30/4	2.56
230	L	I(15) L(75) V(5)T . (1)MFE	0.15	5/1	4.79
158	L	L(64) . (23) R(7)VM K(1) A(1)DCH YSQ	0.24	1/0	4.97

Table 6. The top 25% of residues in 1oihA at the interface with 1oihB. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 7.		
res	type	disruptive mutations
221	G	(R) (H) (FW) (K)
226	D	(R) (H) (FW) (Y)
230	L	(R) (Y) (H) (T)
158	L	(Y) (R) (H) (T)

Table 7. List of disruptive mutations for the top 25% of residues in 1oihA, that are at the interface with 1oihB.

Figure 7 shows residues in 1oihA colored by their importance, at the interface with 1oihB.

Interface with 1oihD. Table 8 lists the top 25% of residues at the interface with 1oihD. The following table (Table 9) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 8.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
132	G	G(76) . (22)SE	0.07	12/12	3.56
246	R	S(3) R(89) T(2)H V(1)	0.08	44/5	3.02

continued in next column




Fig. 7. Residues in 1oihA, at the interface with 1oihB, colored by their relative importance. 1oihB is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 1oihA.)

Table 8. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
266	A	. (1) FEQ AL A (84) R (1) G (4) V (2) P (2)	0.09	2/2	4.13
269	D	. (3) ETS D (71) C (4) Y (4) V T (3) H N (1) G (5) F (2) S . (3) WEP	0.09	46/19	2.93
107	W	W (71) Y (4) . (22) F (1) L	0.13	4/0	3.83
134	D	N (1) D (67) . (22) FR YE (3) HQ T (1) S (1) A	0.21	1/0	4.98
245	V	HV (48)	0.25	8/8	3.99

continued in next column

Table 8. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
		Y (18) WS A (4) F C (7) I (5) L (7) . (1) M T (3)			

Table 8. The top 25% of residues in 1oihA at the interface with 1oihD. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 9.		
res	type	disruptive mutations
132	G	(R) (K) (FWH) (EQM)
246	R	(D) (T) (Y) (E)
266	A	(R) (Y) (K) (E)
269	D	(R) (K) (H) (FW)
107	W	(K) (E) (Q) (TD)
134	D	(R) (FWH) (K) (Y)
245	V	(K) (R) (E) (Y)

Table 9. List of disruptive mutations for the top 25% of residues in 1oihA, that are at the interface with 1oihD.

Figure 8 shows residues in 1oihA colored by their importance, at the interface with 1oihD.

3 NOTES ON USING TRACE RESULTS

3.1 Coverage

Trace results are commonly expressed in terms of coverage: the residue is important if its “coverage” is small - that is if it belongs to some small top percentage of residues [100% is all of the residues in a chain], according to trace. The ET results are presented in the form of a table, usually limited to top 25% percent of residues (or to some nearby percentage), sorted by the strength of the presumed evolutionary pressure. (I.e., the smaller the coverage, the stronger the pressure on the residue.) Starting from the top of that list, mutating a couple of residues should affect the protein somehow, with the exact effects to be determined experimentally.

3.2 Known substitutions

One of the table columns is “substitutions” - other amino acid types seen at the same position in the alignment. These amino acid types may be interchangeable at that position in the protein, so if one wants to affect the protein by a point mutation, they should be avoided. For example if the substitutions are “RVK” and the original protein has an R at that position, it is advisable to try anything, but RVK. Conversely, when looking for substitutions which will *not* affect the protein, one may try replacing, R with K, or (perhaps more surprisingly), with




Fig. 8. Residues in 1oihA, at the interface with 1oihD, colored by their relative importance. 1oihD is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 1oihA.)

V. The percentage of times the substitution appears in the alignment is given in the immediately following bracket. No percentage is given in the cases when it is smaller than 1%. This is meant to be a rough guide - due to rounding errors these percentages often do not add up to 100%.

3.3 Surface

To detect candidates for novel functional interfaces, first we look for residues that are solvent accessible (according to DSSP program) by at least 10\AA^2 , which is roughly the area needed for one water molecule to come in the contact with the residue. Furthermore, we require that these residues form a “cluster” of residues which have neighbor within 5\AA from any of their heavy atoms.

Note, however, that, if our picture of protein evolution is correct, the neighboring residues which *are not* surface accessible might be equally important in maintaining the interaction specificity - they should not be automatically dropped from consideration when choosing the set for mutagenesis. (Especially if they form a cluster with the surface residues.)

3.4 Number of contacts

Another column worth noting is denoted “noc/bb”; it tells the number of contacts heavy atoms of the residue in question make across the interface, as well as how many of them are realized through the backbone atoms (if all or most contacts are through the backbone, mutation presumably won’t have strong impact). Two heavy atoms are considered to be “in contact” if their centers are closer than 5\AA .

3.5 Annotation

If the residue annotation is available (either from the pdb file or from other sources), another column, with the header “annotation” appears. Annotations carried over from PDB are the following: site

(indicating existence of related site record in PDB), S-S (disulfide bond forming residue), hb (hydrogen bond forming residue, jb (james bond forming residue), and sb (for salt bridge forming residue).

3.6 Mutation suggestions

Mutation suggestions are completely heuristic and based on complementarity with the substitutions found in the alignment. Note that they are meant to be **disruptive** to the interaction of the protein with its ligand. The attempt is made to complement the following properties: small [AVGSTC], medium [LPNQDEMIK], large [WIFYHR], hydrophobic [LPVAMWFI], polar [GTCY]; positively [KHR], or negatively [DE] charged, aromatic [WIFYH], long aliphatic chain [EKRQM], OH-group possession [SDETY], and NH2 group possession [NQRK]. The suggestions are listed according to how different they appear to be from the original amino acid, and they are grouped in round brackets if they appear equally disruptive. From left to right, each bracketed group of amino acid types resembles more strongly the original (i.e. is, presumably, less disruptive) These suggestions are tentative - they might prove disruptive to the fold rather than to the interaction. Many researcher will choose, however, the straightforward alanine mutations, especially in the beginning stages of their investigation.

4 APPENDIX

4.1 File formats

Files with extension “ranks_sorted” are the actual trace results. The fields in the table in this file:

- alignment# number of the position in the alignment
- residue# residue number in the PDB file
- type amino acid type
- rank rank of the position according to older version of ET
- variability has two subfields:
 1. number of different amino acids appearing in in this column of the alignment
 2. their type
- rho ET score - the smaller this value, the lesser variability of this position across the branches of the tree (and, presumably, the greater the importance for the protein)
- cvg coverage - percentage of the residues on the structure which have this rho or smaller
- gaps percentage of gaps in this column

4.2 Color schemes used

The following color scheme is used in figures with residues colored by cluster size: black is a single-residue cluster; clusters composed of more than one residue colored according to this hierarchy (ordered by descending size): red, blue, yellow, green, purple, azure, turquoise, brown, coral, magenta, LightSalmon, SkyBlue, violet, gold, bisque, LightSlateBlue, orchid, RosyBrown, MediumAquamarine, DarkOliveGreen, CornflowerBlue, grey55, burlywood, LimeGreen, tan, DarkOrange, DeepPink, maroon, BlanchedAlmond.

The colors used to distinguish the residues by the estimated evolutionary pressure they experience can be seen in Fig. 9.




Fig. 9. Coloring scheme used to color residues by their relative importance.

4.3 Credits

4.3.1 Alistat *alistat* reads a multiple sequence alignment from the file and shows a number of simple statistics about it. These statistics include the format, the number of sequences, the total number of residues, the average and range of the sequence lengths, and the alignment length (e.g. including gap characters). Also shown are some percent identities. A percent pairwise alignment identity is defined as $(\text{idents} / \text{MIN}(\text{len1}, \text{len2}))$ where *idents* is the number of exact identities and *len1*, *len2* are the unaligned lengths of the two sequences. The "average percent identity", "most related pair", and "most unrelated pair" of the alignment are the average, maximum, and minimum of all $(N)(N-1)/2$ pairs, respectively. The "most distant seq" is calculated by finding the maximum pairwise identity (best relative) for all *N* sequences, then finding the minimum of these *N* numbers (hence, the most outlying sequence). *alistat* is copyrighted by HHMI/Washington University School of Medicine, 1992-2001, and freely distributed under the GNU General Public License.

4.3.2 CE To map ligand binding sites from different source structures, *report_maker* uses the CE program: <http://cl.sdsc.edu/>. Shindyalov IN, Bourne PE (1998) "Protein structure alignment by incremental combinatorial extension (CE) of the optimal path". *Protein Engineering* 11(9) 739-747.

4.3.3 DSSP In this work a residue is considered solvent accessible if the DSSP program finds it exposed to water by at least 10\AA^2 , which is roughly the area needed for one water molecule to come in the contact with the residue. DSSP is copyrighted by W. Kabsch, C. Sander and MPI-MF, 1983, 1985, 1988, 1994 1995, CMBI version by Elmar.Krieger@cmbi.kun.nl November 18,2002,

<http://www.cmbi.kun.nl/gv/dssp/descrip.html>.

4.3.4 HSSP Whenever available, *report_maker* uses HSSP alignment as a starting point for the analysis (sequences shorter than 75% of the query are taken out, however); R. Schneider, A. de Daruvar, and C. Sander. "The HSSP database of protein structure-sequence alignments." *Nucleic Acids Res.*, 25:226-230, 1997.

<http://swift.cmbi.kun.nl/swift/hssp/>

4.3.5 LaTeX The text for this report was processed using L^AT_EX; Leslie Lamport, "LaTeX: A Document Preparation System Addison-Wesley," Reading, Mass. (1986).

4.3.6 Muscle When making alignments "from scratch", *report_maker* uses Muscle alignment program: Edgar, Robert C. (2004), "MUSCLE: multiple sequence alignment with high accuracy and high throughput." *Nucleic Acids Research* 32(5), 1792-97.

<http://www.drive5.com/muscle/>

4.3.7 Pymol The figures in this report were produced using Pymol. The scripts can be found in the attachment. Pymol is an open-source application copyrighted by DeLano Scientific LLC (2005). For more information about Pymol see <http://pymol.sourceforge.net/>. (Note for Windows users: the attached package needs to be unzipped for Pymol to read the scripts and launch the viewer.)

4.4 Note about ET Viewer

Dan Morgan from the Lichtarge lab has developed a visualization tool specifically for viewing trace results. If you are interested, please visit:

<http://mammoth.bcm.tmc.edu/traceview/>

The viewer is self-unpacking and self-installing. Input files to be used with ETV (extension .etvx) can be found in the attachment to the main report.

4.5 Citing this work

The method used to rank residues and make predictions in this report can be found in Mihalek, I., I. Reš, O. Lichtarge. (2004). "A Family of Evolution-Entropy Hybrid Methods for Ranking of Protein Residues by Importance" *J. Mol. Bio.* **336**: 1265-82. For the original version of ET see O. Lichtarge, H.Bourne and F. Cohen (1996). "An Evolutionary Trace Method Defines Binding Surfaces Common to Protein Families" *J. Mol. Bio.* **257**: 342-358.

report_maker itself is described in Mihalek I., I. Res and O. Lichtarge (2006). "Evolutionary Trace Report Maker: a new type of service for comparative analysis of proteins." *Bioinformatics* **22**:1656-7.

4.6 About report_maker

report_maker was written in 2006 by Ivana Mihalek. The 1D ranking visualization program was written by Ivica Reš. *report_maker* is copyrighted by Lichtarge Lab, Baylor College of Medicine, Houston.

4.7 Attachments

The following files should accompany this report:

- *loiha.complex.pdb* - coordinates of *loiha* with all of its interacting partners
- *loiha.etvx* - ET viewer input file for *loiha*
- *loiha.cluster_report.summary* - Cluster report summary for *loiha*
- *loiha.ranks* - Ranks file in sequence order for *loiha*
- *loiha.clusters* - Cluster descriptions for *loiha*
- *loiha.msf* - the multiple sequence alignment used for the chain *loiha*

- 1oihA.descr - description of sequences used in 1oihA msf
- 1oihA.ranks_sorted - full listing of residues and their ranking for 1oihA
- 1oihA.1oihCNA1300.if.pml - Pymol script for Figure 5
- 1oihA.cbcvg - used by other 1oihA – related pymol scripts
- 1oihA.1oihANA1301.if.pml - Pymol script for Figure 6
- 1oihA.1oihB.if.pml - Pymol script for Figure 7
- 1oihA.1oihD.if.pml - Pymol script for Figure 8