


2jo8

Evolutionary trace report by **report_maker**

July 7, 2010



4.3.3	DSSP	4
4.3.4	HSSP	4
4.3.5	LaTex	4
4.3.6	Muscle	4
4.3.7	Pymol	4
4.4	Note about ET Viewer	4
4.5	Citing this work	5
4.6	About report_maker	5
4.7	Attachments	5

CONTENTS

1 Introduction

2 Chain 2jo8A

- 2.1 Q5E9L6 overview
- 2.2 Multiple sequence alignment for 2jo8A
- 2.3 Residue ranking in 2jo8A
- 2.4 Top ranking residues in 2jo8A and their position on the structure
 - 2.4.1 Clustering of residues at 25% coverage.
 - 2.4.2 Overlap with known functional surfaces at 25% coverage.

3 Notes on using trace results

- 3.1 Coverage
- 3.2 Known substitutions
- 3.3 Surface
- 3.4 Number of contacts
- 3.5 Annotation
- 3.6 Mutation suggestions

4 Appendix

- 4.1 File formats
- 4.2 Color schemes used
- 4.3 Credits
 - 4.3.1 **Alistat**
 - 4.3.2 **CE**

1 INTRODUCTION

From the original Protein Data Bank entry (PDB id 2jo8):

Title: Solution structure of c-terminal sarah domain, database sterile 20-like kinase 1 (mst1)

Compound: Mol id: 1; molecule: serine/threonine-protein kinase 4; chain: a, b; fragment: c-terminal sarah domain, database residues 432- 480; synonym: ste20-like kinase mst1, mst-1, mammalian ste20- like protein kinase 1, serine/threonine-protein kinase krs- 2; ec: 2.7.11.1; engineered: yes

1 **Organism, scientific name:** Homo Sapiens;

2jo8 contains a single unique chain 2jo8A (51 residues long) and its homologue 2jo8B. This is an NMR-determined structure – in this report the first model in the file was used.

2 CHAIN 2JO8A

2.1 Q5E9L6 overview

2 From SwissProt, id Q5E9L6, 100% identical to 2jo8A:

Description: Serine/threonine kinase 4.

3 **Organism, scientific name:** Bos taurus (Bovine).

3 **Taxonomy:** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Laurasiatheria; Cetartiodactyla; Ruminantia; Pecora; Bovidae; Bovinae; Bos.

2.2 Multiple sequence alignment for 2jo8A

4 For the chain 2jo8A, the alignment 2jo8A.msf (attached) with 25 sequences was used. The alignment was downloaded from the HSSP database, and fragments shorter than 75% of the query as well as duplicate sequences were removed. It can be found in the attachment to this report, under the name of 2jo8A.msf. Its statistics, from the *alistat* program are the following:



Fig. 1. Residues 1-51 in 2jo8A colored by their relative importance. (See Appendix, Fig.5, for the coloring scheme.)

```

Format:                MSF
Number of sequences:  25
Total number of residues: 1217
Smallest:             41
Largest:              51
Average length:       48.7
Alignment length:     51
Average identity:     56%
Most related pair:    98%
Most unrelated pair:  34%
Most distant seq:     49%
  
```

Furthermore, 11% of residues show as conserved in this alignment. The alignment consists of 16% eukaryotic (8% vertebrata, 4% arthropoda) sequences. (Descriptions of some sequences were not readily available.) The file containing the sequence descriptions can be found in the attachment, under the name 2jo8A.descr.

2.3 Residue ranking in 2jo8A

The 2jo8A sequence is shown in Fig. 1, with each residue colored according to its estimated importance. The full listing of residues in 2jo8A can be found in the file called 2jo8A.ranks_sorted in the attachment.

2.4 Top ranking residues in 2jo8A and their position on the structure

In the following we consider residues ranking among top 25% of residues in the protein . Figure 2 shows residues in 2jo8A colored by their importance: bright red and yellow indicate more conserved/important residues (see Appendix for the coloring scheme). A Pymol script for producing this figure can be found in the attachment.

2.4.1 Clustering of residues at 25% coverage. Fig. 3 shows the top 25% of all residues, this time colored according to clusters they belong to. The clusters in Fig.3 are composed of the residues listed in Table 1.

Table 1.		
cluster color	size	member residues
red	7	37, 40, 41, 43, 44, 47, 51
blue	4	23, 26, 27, 29
yellow	2	7, 15

Table 1. Clusters of top ranking residues in 2jo8A.

2.4.2 Overlap with known functional surfaces at 25% coverage. The name of the ligand is composed of the source PDB identifier and the heteroatom name used in that file.




Fig. 2. Residues in 2jo8A, colored by their relative importance. Clockwise: front, back, top and bottom views.




Fig. 3. Residues in 2jo8A, colored according to the cluster they belong to: red, followed by blue and yellow are the largest clusters (see Appendix for the coloring scheme). Clockwise: front, back, top and bottom views. The corresponding Pymol script is attached.

Interface with 2jo8B. Table 2 lists the top 25% of residues at the interface with 2jo8B. The following table (Table 3) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
37	Y	Y(100)	0.12	284/22	1.66
40	K	K(100)	0.12	60/10	2.80
41	R	R(100)	0.12	215/16	2.28
43	P	P(100)	0.12	74/16	2.61
44	I	I(100)	0.12	379/35	2.14
51	K	K(100)	0.12	163/0	1.82
26	M	M(95) K(4)	0.14	123/31	2.53
29	E	E(95) D(4)	0.16	136/16	1.59
7	L	L(92) V(4) .(4)	0.18	194/17	2.14
15	L	L(95) V(4)	0.20	264/15	2.06
47	A	A(95) V(4)	0.22	166/40	2.34

Table 2. The top 25% of residues in 2jo8A at the interface with 2jo8B. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

res	type	disruptive mutations
37	Y	(K) (QM) (NEVLAPIR) (D)
40	K	(Y) (FTW) (SVCAG) (HD)
41	R	(TD) (SYEVCLAPIG) (FMW) (N)
43	P	(YR) (TH) (SKECG) (FQWD)
44	I	(YR) (TH) (SKECG) (FQWD)
51	K	(Y) (FTW) (SVCAG) (HD)
26	M	(Y) (T) (H) (SCG)
29	E	(FWH) (R) (YVCAG) (T)
7	L	(Y) (R) (H) (T)
15	L	(YR) (H) (TKE) (SQCDG)
47	A	(KYER) (QHD) (N) (FTMW)

Table 3. List of disruptive mutations for the top 25% of residues in 2jo8A, that are at the interface with 2jo8B.

Figure 4 shows residues in 2jo8A colored by their importance, at the interface with 2jo8B.

3 NOTES ON USING TRACE RESULTS

3.1 Coverage

Trace results are commonly expressed in terms of coverage: the residue is important if its “coverage” is small - that is if it belongs to some small top percentage of residues [100% is all of the residues in a chain], according to trace. The ET results are presented in the form of a table, usually limited to top 25% percent of residues (or to some nearby percentage), sorted by the strength of the presumed




Fig. 4. Residues in 2jo8A, at the interface with 2jo8B, colored by their relative importance. 2jo8B is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 2jo8A.)

evolutionary pressure. (I.e., the smaller the coverage, the stronger the pressure on the residue.) Starting from the top of that list, mutating a couple of residues should affect the protein somehow, with the exact effects to be determined experimentally.

3.2 Known substitutions

One of the table columns is “substitutions” - other amino acid types seen at the same position in the alignment. These amino acid types may be interchangeable at that position in the protein, so if one wants to affect the protein by a point mutation, they should be avoided. For example if the substitutions are “RVK” and the original protein has an R at that position, it is advisable to try anything, but RVK. Conversely, when looking for substitutions which will *not* affect the protein, one may try replacing, R with K, or (perhaps more surprisingly), with V. The percentage of times the substitution appears in the alignment is given in the immediately following bracket. No percentage is given in the cases when it is smaller than 1%. This is meant to be a rough guide - due to rounding errors these percentages often do not add up to 100%.

3.3 Surface

To detect candidates for novel functional interfaces, first we look for residues that are solvent accessible (according to DSSP program) by at least 10\AA^2 , which is roughly the area needed for one water molecule to come in the contact with the residue. Furthermore, we require that these residues form a “cluster” of residues which have neighbor within 5\AA from any of their heavy atoms.

Note, however, that, if our picture of protein evolution is correct, the neighboring residues which *are not* surface accessible might be equally important in maintaining the interaction specificity - they

which is roughly the area needed for one water molecule to come in the contact with the residue. DSSP is copyrighted by W. Kabsch, C. Sander and MPI-MF, 1983, 1985, 1988, 1994 1995, CMBI version by Elmar.Krieger@cmbi.kun.nl November 18,2002,

<http://www.cmbi.kun.nl/gv/dssp/descrip.html>.

4.3.4 HSSP Whenever available, report_maker uses HSSP alignment as a starting point for the analysis (sequences shorter than 75% of the query are taken out, however); R. Schneider, A. de Daruvar, and C. Sander. "The HSSP database of protein structure-sequence alignments." *Nucleic Acids Res.*, 25:226–230, 1997.

<http://swift.cmbi.kun.nl/swift/hssp/>

4.3.5 LaTeX The text for this report was processed using L^AT_EX; Leslie Lamport, "LaTeX: A Document Preparation System Addison-Wesley," Reading, Mass. (1986).

4.3.6 Muscle When making alignments "from scratch", report_maker uses Muscle alignment program: Edgar, Robert C. (2004), "MUSCLE: multiple sequence alignment with high accuracy and high throughput." *Nucleic Acids Research* 32(5), 1792-97.

<http://www.drive5.com/muscle/>

4.3.7 Pymol The figures in this report were produced using Pymol. The scripts can be found in the attachment. Pymol is an open-source application copyrighted by DeLano Scientific LLC (2005). For more information about Pymol see <http://pymol.sourceforge.net/>. (Note for Windows users: the attached package needs to be unzipped for Pymol to read the scripts and launch the viewer.)

4.4 Note about ET Viewer

Dan Morgan from the Lichtarge lab has developed a visualization tool specifically for viewing trace results. If you are interested, please visit:

<http://mammoth.bcm.tmc.edu/traceview/>

The viewer is self-unpacking and self-installing. Input files to be used with ETV (extension .etvx) can be found in the attachment to the main report.

4.5 Citing this work

The method used to rank residues and make predictions in this report can be found in Mihalek, I., I. Reš, O. Lichtarge. (2004). "A Family of Evolution-Entropy Hybrid Methods for Ranking of Protein Residues by Importance" *J. Mol. Bio.* **336**: 1265-82. For the original version of ET see O. Lichtarge, H.Bourne and F. Cohen (1996). "An Evolutionary Trace Method Defines Binding Surfaces Common to Protein Families" *J. Mol. Bio.* **257**: 342-358.

report_maker itself is described in Mihalek I., I. Res and O. Lichtarge (2006). "Evolutionary Trace Report Maker: a new type of service for comparative analysis of proteins." *Bioinformatics* **22**:1656-7.

4.6 About report_maker

report_maker was written in 2006 by Ivana Mihalek. The 1D ranking visualization program was written by Ivica Reš. report_maker is copyrighted by Lichtarge Lab, Baylor College of Medicine, Houston.

4.7 Attachments

The following files should accompany this report:

- 2jo8A.complex.pdb - coordinates of 2jo8A with all of its interacting partners
- 2jo8A.etvx - ET viewer input file for 2jo8A
- 2jo8A.cluster_report.summary - Cluster report summary for 2jo8A
- 2jo8A.ranks - Ranks file in sequence order for 2jo8A
- 2jo8A.clusters - Cluster descriptions for 2jo8A
- 2jo8A.msf - the multiple sequence alignment used for the chain 2jo8A
- 2jo8A.descr - description of sequences used in 2jo8A msf
- 2jo8A.ranks_sorted - full listing of residues and their ranking for 2jo8A
- 2jo8A.2jo8B.if.pml - Pymol script for Figure 4
- 2jo8A.cbvcg - used by other 2jo8A – related pymol scripts