


3ecs

Evolutionary trace report by **report_maker**

August 4, 2010



CONTENTS

1 Introduction

2 Chain 3ecsC

- 2.1 Q14232 overview
- 2.2 Multiple sequence alignment for 3ecsC
- 2.3 Residue ranking in 3ecsC
- 2.4 Top ranking residues in 3ecsC and their position on the structure
 - 2.4.1 Clustering of residues at 25% coverage.
 - 2.4.2 Overlap with known functional surfaces at 25% coverage.
 - 2.4.3 Possible novel functional surfaces at 25% coverage.

3 Notes on using trace results

- 3.1 Coverage
- 3.2 Known substitutions
- 3.3 Surface
- 3.4 Number of contacts
- 3.5 Annotation
- 3.6 Mutation suggestions

4 Appendix

- 4.1 File formats
- 4.2 Color schemes used
- 4.3 Credits

4.3.1	Alistat	11
4.3.2	CE	11
4.3.3	DSSP	11
4.3.4	HSSP	11
4.3.5	LaTex	11
4.3.6	Muscle	11
4.3.7	Pymol	11
4.4	Note about ET Viewer	11
4.5	Citing this work	11
4.6	About report_maker	11
4.7	Attachments	11

1 INTRODUCTION

From the original Protein Data Bank entry (PDB id 3ecs):

Title: Crystal structure of human eif2b alpha

Compound: Mol id: 1; molecule: translation initiation factor eif-2b subunit alpha; chain: a, b, c, d, e, f, g, h; synonym: eukaryotic translation initiation factor 2b alpha, eif-2b gdp-gtp exchange factor subunit alpha; engineered: yes

Organism, scientific name: Homo Sapiens;

3ecs contains a single unique chain 3ecsC (294 residues long) and its homologues 3ecsF, 3ecsA, 3ecsE, 3ecsB, 3ecsH, 3ecsD, and 3ecsG.

1

1 2 CHAIN 3ECSC

1

2.1 Q14232 overview

1

From SwissProt, id Q14232, 85% identical to 3ecsC:

Description: Translation initiation factor eIF-2B alpha subunit (eIF-2B GDP-GTP exchange factor).

Organism, scientific name: Homo sapiens (Human).

Taxonomy: Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Catarrhini; Hominidae; Homo.

Function: Catalyzes the exchange of eukaryotic initiation factor 2-bound GDP for GTP.

Subunit: Complex of five different subunits; alpha, beta, gamma, delta and epsilon.

Interaction:

Disease: Defects in EIF2B1 are a cause of leukoencephalopathy with vanishing white matter (VWM) [MIM:603896]. VWM is a brain disease that occurs mainly in children. The neurological signs of VWM include progressive cerebellar ataxia, spasticity, inconstant optic atrophy and relatively preserved mental abilities. The disease is chronic-progressive with, in most individuals, additional episodes of rapid deterioration following febrile infections or minor head trauma.

Head trauma leads only to motor deterioration, whereas infections with fever may end in coma. Death occurs after a variable period of

11




Fig. 1. Residues 1-150 in 3ecsC colored by their relative importance. (See Appendix, Fig.12, for the coloring scheme.)

a few years to a few decades, usually following an episode of fever and coma.

Similarity: Belongs to the EIF-2B alpha/beta/delta subunits family.

About: This Swiss-Prot entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use as long as its content is in no way modified and this statement is not removed.

2.2 Multiple sequence alignment for 3ecsC

For the chain 3ecsC, the alignment 3ecsC.msf (attached) with 570 sequences was used. The alignment was downloaded from the HSSP database, and fragments shorter than 75% of the query as well as duplicate sequences were removed. It can be found in the attachment to this report, under the name of 3ecsC.msf. Its statistics, from the *alistat* program are the following:

```

Format:                MSF
Number of sequences:   570
Total number of residues: 93711
Smallest:              32
Largest:               294
Average length:        164.4
Alignment length:      294
Average identity:       34%
Most related pair:     99%
Most unrelated pair:   0%
Most distant seq:      31%

```

Furthermore, <1% of residues show as conserved in this alignment.

The alignment consists of 9% eukaryotic (3% vertebrata, <1% arthropoda, 3% fungi, <1% plantae), 1% prokaryotic, and 4% archaean sequences. (Descriptions of some sequences were not readily available.) The file containing the sequence descriptions can be found in the attachment, under the name 3ecsC.descr.

2.3 Residue ranking in 3ecsC

The 3ecsC sequence is shown in Figs. 1–2, with each residue colored according to its estimated importance. The full listing of residues in 3ecsC can be found in the file called 3ecsC.ranks_sorted in the attachment.




Fig. 2. Residues 151-307 in 3ecsC colored by their relative importance. (See Appendix, Fig.12, for the coloring scheme.)

2.4 Top ranking residues in 3ecsC and their position on the structure

In the following we consider residues ranking among top 25% of residues in the protein. Figure 3 shows residues in 3ecsC colored by their importance: bright red and yellow indicate more conserved/important residues (see Appendix for the coloring scheme). A Pymol script for producing this figure can be found in the attachment.




Fig. 3. Residues in 3ecsC, colored by their relative importance. Clockwise: front, back, top and bottom views.

2.4.1 Clustering of residues at 25% coverage. Fig. 4 shows the top 25% of all residues, this time colored according to clusters they belong to. The clusters in Fig.4 are composed of the residues listed in Table 1.

Table 1.		
cluster color	size	member residues
red	73	111, 118, 125, 126, 127, 128, 129, 131, 132, 133, 134, 137, 138, 141

continued in next column




Fig. 4. Residues in 3ecsC, colored according to the cluster they belong to: red, followed by blue and yellow are the largest clusters (see Appendix for the coloring scheme). Clockwise: front, back, top and bottom views. The corresponding Pymol script is attached.

Table 1. continued		
cluster color	size	member residues
		145, 150, 152, 153, 154, 155, 156 157, 160, 161, 164, 165, 168, 173 175, 177, 178, 180, 181, 182, 183 184, 190, 191, 193, 195, 196, 197 198, 199, 200, 201, 202, 203, 204 205, 206, 207, 208, 209, 210, 211 212, 213, 215, 216, 217, 219, 224 225, 226, 227, 228, 229, 230, 231 233, 234, 235

Table 1. Clusters of top ranking residues in 3ecsC.

2.4.2 Overlap with known functional surfaces at 25% coverage.

The name of the ligand is composed of the source PDB identifier and the heteroatom name used in that file.

Interface with 3ecsE. Table 2 lists the top 25% of residues at the interface with 3ecsE. The following table (Table 3) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 2.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
203	N	N (71) D (9) EY S (13)	0.08	4/3	4.60

continued in next column

Table 2. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
202	E	T (1) HQK CAIRG. E (21) A (50) K (2) L (3) T (1) R (4) S (8) NIG WMQV (2) .DPC	0.19	33/2	2.81

Table 2. The top 25% of residues in 3ecsC at the interface with 3ecsE. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 3.		
res	type	disruptive mutations
203	N	(Y) (FW) (H) (T)
202	E	(H) (FW) (Y) (R)

Table 3. List of disruptive mutations for the top 25% of residues in 3ecsC, that are at the interface with 3ecsE.




Fig. 5. Residues in 3ecsC, at the interface with 3ecsE, colored by their relative importance. 3ecsE is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 3ecsC.)

Figure 5 shows residues in 3ecsC colored by their importance, at the interface with 3ecsE.

Sulfate ion binding site. By analogy with 3ecsF – 3ecsFSO4506 interface. Table 4 lists the top 25% of residues at the interface with 3ecsFSO4506 (sulfate ion). The following table (Table 5) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 4.						
res	type	subst's (%)	cvg	noc/ bb	dist (Å)	antn
196	G	G(97)SA P.N	0.01	3/3	4.77	
198	E	E(17) H(10) D(66) Q(1)A K(1)YSN .R	0.02	1/1	4.97	
197	A	A(83) T(5)M C(5) S(2).LV I	0.05	5/5	3.77	
134	V	V(35) L(6) A(53) I(2)YKM .SG	0.06	1/1	4.82	
132	R	R(22) S(10) G(51)V T(2) H(1) K(5) E(1)DL N(1)AMQ .P	0.11	25/10	3.07	site
131	S	S(45) A(13) Y(20) T(9) W(3)V H(1)GNE F(2)LRI .	0.13	18/10	2.83	site
234	K	K(48) T(27) . (18) S(5)HDE	0.13	6/0	3.56	
133	V	V(19) T(58) A(9) L(4)GDH	0.14	7/4	3.34	site

continued in next column

Table 4. continued						
res	type	subst's (%)	cvg	noc/ bb	dist (Å)	antn
		S(2)QFW I(1)R.Y CN				

Table 4. The top 25% of residues in 3ecsC at the interface with sulfate ion.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 5.		
res	type	disruptive mutations
196	G	(R)(KE)(H)(FW)
198	E	(FW)(H)(Y)(VCAG)
197	A	(R)(K)(Y)(E)
134	V	(R)(Y)(K)(E)
132	R	(Y)(T)(D)(E)
131	S	(K)(R)(Q)(H)
234	K	(Y)(FW)(T)(VA)
133	V	(KER)(Y)(Q)(D)

Table 5. List of disruptive mutations for the top 25% of residues in 3ecsC, that are at the interface with sulfate ion.




Fig. 6. Residues in 3ecsC, at the interface with sulfate ion, colored by their relative importance. The ligand (sulfate ion) is colored green. Atoms further than 30Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain 3ecsC.)

Figure 6 shows residues in 3ecsC colored by their importance, at the interface with 3ecsFSO4506.

Interface with 3ecsF. By analogy with 3ecsD – 3ecsF interface. Table 6 lists the top 25% of residues at the interface with 3ecsF. The following table (Table 7) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 6.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
203	N	N (71) D (9) EY S (13) T (1) HQK CAIRG.	0.08	11/4	3.53
202	E	E (21) A (50) K (2) L (3) T (1) R (4) S (8) NIG WMQV (2) .DPC	0.19	42/7	3.12




Fig. 7. Residues in 3ecsC, at the interface with 3ecsF, colored by their relative importance. 3ecsF is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 3ecsC.)

Table 6. The top 25% of residues in 3ecsC at the interface with 3ecsF. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 7.		
res	type	disruptive mutations
203	N	(Y) (FW) (H) (T)
202	E	(H) (FW) (Y) (R)

Table 7. List of disruptive mutations for the top 25% of residues in 3ecsC, that are at the interface with 3ecsF.

Figure 7 shows residues in 3ecsC colored by their importance, at the interface with 3ecsF.

Interface with 3ecsA. Table 8 lists the top 25% of residues at the interface with 3ecsA. The following table (Table 9) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 8.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
211	G	G (99) .	0.00	10/10	2.94
180	D	D (85) I (5) L (1) E (4) PA V (1) NT .	0.02	20/8	2.96

continued in next column

Table 8. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
154	E	CM E (83) D (8) S (1) LT Q (1) N (1) HI . VMGR	0.03	3/0	4.53
157	P	P (93) A (1) RE T (1) D (1) S . I G	0.03	5/2	3.81
156	Q	R (78) A (7) S (2) HP Q (3) Y (1) C G (2) NFE K . LD	0.09	26/5	3.38
150	V	V (77) T I (4) A (10) G L (3) F C (1) SM . H	0.12	1/1	4.70
182	A	A (58) M (26)	0.12	15/12	3.35

continued in next column

Table 8. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
213	N	Q(1)E G(2) L(1) S(4)K T(3).CV N Y(58) H(2) F(4) L(4) S(7) A(9)T R(1) M(3)G N(4)CK. V	0.15	15/0	3.35
184	G	A(33) L(1) G(36) S(7) V(3) R(5) F(5) P(1) Y(1)TWD E.HIK	0.16	23/23	3.08
210	I	L(3)C V(17) I(64) A(4) N(1) T(2) M(4) S(1).K	0.16	30/8	3.66
177	V	I(4) V(22) L(52) Y(9)Q T(4)PF A(1)C M(1)S.N K	0.17	36/30	2.94
217	V	T(1) L(32) A(1) I(18) M(5) V(35)EQ N(1)H C(1)RS.	0.20	14/2	3.67
178	V	V(16) I(68)	0.21	6/2	4.17

continued in next column

Table 8. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
175	V	A(2)C L(4) M(1) N(1)DQS EGY. T(1)H V(51)P C(10) F(3) A(8) T(11) H(4) Y(4)L S(1)NM I(1)QG.	0.22	36/33	2.48
181	A	A(20) S(40) G(6) N(23) H(1)LV T(4) P(1)F.D C	0.24	54/41	3.07
183	V	M(4) V(31) A(38) I(7) C(1) S(3) L(5) T(2)QP G(1)F H(2)N.	0.24	20/11	2.49

Table 8. The top 25% of residues in 3ecsC at the interface with 3ecsA. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/ bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 9.		
res	type	disruptive mutations
211	G	(KER)(FQMWHD)(NLPI)(Y)
180	D	(R)(H)(FW)(Y)
154	E	(H)(FW)(Y)(R)
157	P	(R)(Y)(H)(K)
156	Q	(Y)(H)(FTW)(CG)
150	V	(R)(K)(E)(Y)
182	A	(Y)(R)(H)(KE)
213	N	(Y)(H)(E)(FW)

continued in next column

res	type	disruptive mutations
184	G	(R) (K) (E) (Q)
210	I	(Y) (R) (H) (T)
177	V	(R) (Y) (E) (K)
217	V	(Y) (R) (E) (K)
178	V	(R) (KY) (E) (H)
175	V	(R) (KE) (Y) (H)
181	A	(R) (K) (E) (Y)
183	V	(R) (Y) (E) (K)

Table 9. List of disruptive mutations for the top 25% of residues in 3ecsC, that are at the interface with 3ecsA.




Fig. 8. Residues in 3ecsC, at the interface with 3ecsA, colored by their relative importance. 3ecsA is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 3ecsC.)

Figure 8 shows residues in 3ecsC colored by their importance, at the interface with 3ecsA.

Sulfate ion binding site. Table 10 lists the top 25% of residues at the interface with 3ecsCSO4503 (sulfate ion). The following table (Table 11) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)	antn
196	G	G(97)SA P.N	0.01	2/2	4.54	
197	A	A(83)	0.05	3/3	4.40	

continued in next column

res	type	subst's (%)	cvg	noc/ bb	dist (Å)	antn
231	E	T(5)M C(5) S(2).LV I E(35) P(52) S(1) Q(1) G(4) D(1) . (2)TAH	0.05	4/0	4.33	
132	R	R(22) S(10) G(51)V T(2) H(1) K(5) E(1)DL N(1)AMQ .P	0.11	35/14	3.07	site
131	S	S(45) A(13) Y(20) T(9) W(3)V H(1)GNE F(2)LRI .	0.13	17/10	2.94	site
234	K	K(48) T(27) . (18) S(5)HDE	0.13	6/0	2.83	
133	V	V(19) T(58) A(9) L(4)GDH S(2)QFW I(1)R.Y CN	0.14	10/5	3.15	site

Table 10. The top 25% of residues in 3ecsC at the interface with sulfate ion.(Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

res	type	disruptive mutations
196	G	(R) (KE) (H) (FW)
197	A	(R) (K) (Y) (E)
231	E	(H) (FW) (R) (Y)

continued in next column

res	type	disruptive mutations
132	R	(Y) (T) (D) (E)
131	S	(K) (R) (Q) (H)
234	K	(Y) (FW) (T) (VA)
133	V	(KER) (Y) (Q) (D)

Table 11. List of disruptive mutations for the top 25% of residues in 3ecsC, that are at the interface with sulfate ion.




Fig. 10. A possible active surface on the chain 3ecsC. The larger cluster it belongs to is shown in blue.




Fig. 9. Residues in 3ecsC, at the interface with sulfate ion, colored by their relative importance. The ligand (sulfate ion) is colored green. Atoms further than 30 Å away from the geometric center of the ligand, as well as on the line of sight to the ligand were removed. (See Appendix for the coloring scheme for the protein chain 3ecsC.)

Figure 9 shows residues in 3ecsC colored by their importance, at the interface with 3ecsCSO4503.

2.4.3 Possible novel functional surfaces at 25% coverage. One group of residues is conserved on the 3ecsC surface, away from (or substantially larger than) other functional sites and interfaces recognizable in PDB entry 3ecs. It is shown in Fig. 10. The right panel shows (in blue) the rest of the larger cluster this surface belongs to. The residues belonging to this surface "patch" are listed in Table 12, while Table 13 suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	substitutions(%)	cvg
225	P	P(87)AR(3)NW(1) K(2)SGDE.Q	0.04

continued in next column

res	type	substitutions(%)	cvg
226	F	F(73)L(4)V(17)I YM(1)T(1).C	0.08
227	Y	Y(64)I(7)TL(8) M(6)F(5)V(4)WH. A	0.12
224	K	K(21)V(31)I(40) T(3)L(1)CR.AM	0.17
191	D	D(61)S(5)Q(4) H(2)N(11)T(8) A(1)E(2)GRM.K	0.18
118	F	F(21)H(8)TL(34) K(6)M(1)R(5)E C(2)V(3)YA.(8)D I(1)GQWSP	0.20
122	G	N(18)G(57)D(6) R(1)K(2)H(1) Q(1)E(1)T.(6) S(1)PLVAC	0.25

Table 12. Residues forming surface "patch" in 3ecsC.

res	type	disruptive mutations
225	P	(Y) (R) (H) (T)
226	F	(K) (E) (Q) (R)
227	Y	(K) (Q) (E) (R)
224	K	(Y) (FW) (T) (H)
191	D	(R) (FW) (H) (Y)
118	F	(K) (E) (T) (D)
122	G	(R) (E) (K) (H)

Table 13. Disruptive mutations for the surface patch in 3ecsC.

Another group of surface residues is shown in Fig. 11. The right panel shows (in blue) the rest of the larger cluster this surface belongs to. The residues belonging to this surface "patch" are listed in Table 14, while Table 15 suggests possible disruptive replacements for these residues (see Section 3.6).




Fig. 11. Another possible active surface on the chain 3ecsC. The larger cluster it belongs to is shown in blue.

Table 14.				
res	type	substitutions(%)	cvg	antn
211	G	G(99) .	0.00	
196	G	G(97)SAP.N	0.01	
180	D	D(85)I(5)L(1) E(4)PAV(1)NT.CM	0.02	
198	E	E(17)H(10)D(66) Q(1)AK(1)YSN.R	0.02	
154	E	E(83)D(8)S(1)LT Q(1)N(1)HI.VMGR	0.03	
157	P	P(93)A(1)RET(1) D(1)S.IG	0.03	
208	N	N(78)A(7)G(1) S(11)HV.	0.03	
197	A	A(83)T(5)MC(5) S(2).LVI	0.05	
231	E	E(35)P(52)S(1) Q(1)G(4)D(1) . (2)TAH	0.05	
209	K	K(68)P(2)R(15) A(1)V(2)M(2)G S(1)H(1)L(1)IQY E.ND	0.06	
156	Q	R(78)A(7)S(2)HP Q(3)Y(1)CG(2)NF EK.LD	0.09	
160	S	S(5)L(2)Q(64) E(17)GT(3)A(1)R DHNIY.PKF	0.10	
132	R	R(22)S(10)G(51) VT(2)H(1)K(5) E(1)DLN(1)AMQ.P	0.11	site
150	V	V(77)TI(4)A(10) GL(3)FC(1)SM.H	0.12	
182	A	A(58)M(26)Q(1)E G(2)L(1)S(4)K T(3).CVN	0.12	
128	H	H(54)T(17)I(5) L(3)F(3)N(4) Y(6)V(1)AQGDC.M R	0.13	
131	S	S(45)A(13)Y(20) T(9)W(3)VH(1)GN EF(2)LRI.	0.13	site
234	K	K(48)T(27).(18) S(5)HDE	0.13	
213	N	Y(58)H(2)F(4) L(4)S(7)A(9)T R(1)M(3)GN(4)CK .V	0.15	
184	G	A(33)L(1)G(36) S(7)V(3)R(5)	0.16	

continued in next column

res	type	substitutions(%)	cvg	antn
210	I	F(5)P(1)Y(1)TWD E.HIK L(3)CV(17)I(64) A(4)N(1)T(2) M(4)S(1).K	0.16	
129	A	G(24)DC(28) A(28)S(14)TV(1) QENIK.	0.17	
177	V	I(4)V(22)L(52) Y(9)QT(4)PFA(1) CM(1)S.NK	0.17	
165	A	A(59)T(14)L(5) V(7)S(2)I(2) Q(1)GY(3)HDCMK. RNEF	0.20	
217	V	T(1)L(32)A(1) I(18)M(5)V(35)E QN(1)HC(1)RS.	0.20	
173	V	I(68)V(20)L(6) F(2)QMAH.YKTN	0.21	
178	V	V(16)I(68)A(2)C L(4)M(1)N(1)DQS EGY.T(1)H	0.21	
175	V	V(51)PC(10)F(3) A(8)T(11)H(4) Y(4)LS(1)NMI(1) QG.	0.22	
233	F	Y(13)F(14)A S(23)P(1).(17) I(2)EH(11)T(8)C NGL(3)V(1)DQK	0.23	
181	A	A(20)S(40)G(6) N(23)H(1)LV(4) P(1)F.DC	0.24	
183	V	M(4)V(31)A(38) I(7)C(1)S(3) L(5)T(2)QPG(1)F H(2)N.	0.24	

Table 14. Residues forming surface "patch" in 3ecsC.

res	type	disruptive mutations
211	G	(KER)(FQMWHD)(NLPI)(Y)
196	G	(R)(KE)(H)(FW)
180	D	(R)(H)(FW)(Y)
198	E	(FW)(H)(Y)(VCAG)
154	E	(H)(FW)(Y)(R)
157	P	(R)(Y)(H)(K)
208	N	(Y)(H)(E)(FW)
197	A	(R)(K)(Y)(E)
231	E	(H)(FW)(R)(Y)

continued in next column

res	type	disruptive mutations
209	K	(Y)(T)(FW)(CG)
156	Q	(Y)(H)(FTW)(CG)
160	S	(R)(K)(H)(FW)
132	R	(Y)(T)(D)(E)
150	V	(R)(K)(E)(Y)
182	A	(Y)(R)(H)(KE)
128	H	(E)(D)(TQ)(KM)
131	S	(K)(R)(Q)(H)
234	K	(Y)(FW)(T)(VA)
213	N	(Y)(H)(E)(FW)
184	G	(R)(K)(E)(Q)
210	I	(Y)(R)(H)(T)
129	A	(Y)(R)(H)(K)
177	V	(R)(Y)(E)(K)
165	A	(YR)(K)(E)(H)
217	V	(Y)(R)(E)(K)
173	V	(Y)(E)(R)(K)
178	V	(R)(KY)(E)(H)
175	V	(R)(KE)(Y)(H)
233	F	(K)(E)(R)(Q)
181	A	(R)(K)(E)(Y)
183	V	(R)(Y)(E)(K)

Table 15. Disruptive mutations for the surface patch in 3ecsC.

3 NOTES ON USING TRACE RESULTS

3.1 Coverage

Trace results are commonly expressed in terms of coverage: the residue is important if its "coverage" is small - that is if it belongs to some small top percentage of residues [100% is all of the residues in a chain], according to trace. The ET results are presented in the form of a table, usually limited to top 25% percent of residues (or to some nearby percentage), sorted by the strength of the presumed evolutionary pressure. (I.e., the smaller the coverage, the stronger the pressure on the residue.) Starting from the top of that list, mutating a couple of residues should affect the protein somehow, with the exact effects to be determined experimentally.

3.2 Known substitutions

One of the table columns is "substitutions" - other amino acid types seen at the same position in the alignment. These amino acid types may be interchangeable at that position in the protein, so if one wants to affect the protein by a point mutation, they should be avoided. For example if the substitutions are "RVK" and the original protein has an R at that position, it is advisable to try anything, but RVK. Conversely, when looking for substitutions which will *not* affect the protein, one may try replacing, R with K, or (perhaps more surprisingly), with V. The percentage of times the substitution appears in the alignment is given in the immediately following bracket. No percentage is given in the cases when it is smaller than 1%. This is meant to be a rough guide - due to rounding errors these percentages often do not add up to 100%.

3.3 Surface

To detect candidates for novel functional interfaces, first we look for residues that are solvent accessible (according to DSSP program) by at least 10\AA^2 , which is roughly the area needed for one water molecule to come in the contact with the residue. Furthermore, we require that these residues form a “cluster” of residues which have neighbor within 5\AA from any of their heavy atoms.

Note, however, that, if our picture of protein evolution is correct, the neighboring residues which *are not* surface accessible might be equally important in maintaining the interaction specificity - they should not be automatically dropped from consideration when choosing the set for mutagenesis. (Especially if they form a cluster with the surface residues.)

3.4 Number of contacts

Another column worth noting is denoted “noc/bb”; it tells the number of contacts heavy atoms of the residue in question make across the interface, as well as how many of them are realized through the backbone atoms (if all or most contacts are through the backbone, mutation presumably won’t have strong impact). Two heavy atoms are considered to be “in contact” if their centers are closer than 5\AA .

3.5 Annotation

If the residue annotation is available (either from the pdb file or from other sources), another column, with the header “annotation” appears. Annotations carried over from PDB are the following: site (indicating existence of related site record in PDB), S-S (disulfide bond forming residue), hb (hydrogen bond forming residue, jb (james bond forming residue), and sb (for salt bridge forming residue).

3.6 Mutation suggestions

Mutation suggestions are completely heuristic and based on complementarity with the substitutions found in the alignment. Note that they are meant to be **disruptive** to the interaction of the protein with its ligand. The attempt is made to complement the following properties: small [AVGSTC], medium [LPNQDEMILK], large [WPHYHR], hydrophobic [LPVAMWFI], polar [GTCY]; positively [KHR], or negatively [DE] charged, aromatic [WPHYH], long aliphatic chain [EK RQM], OH-group possession [SDETY], and NH2 group possession [NQRK]. The suggestions are listed according to how different they appear to be from the original amino acid, and they are grouped in round brackets if they appear equally disruptive. From left to right, each bracketed group of amino acid types resembles more strongly the original (i.e. is, presumably, less disruptive) These suggestions are tentative - they might prove disruptive to the fold rather than to the interaction. Many researcher will choose, however, the straightforward alanine mutations, especially in the beginning stages of their investigation.

4 APPENDIX

4.1 File formats

Files with extension “ranks_sorted” are the actual trace results. The fields in the table in this file:

- alignment# number of the position in the alignment
- residue# residue number in the PDB file
- type amino acid type
- rank rank of the position according to older version of ET




Fig. 12. Coloring scheme used to color residues by their relative importance.

- variability has two subfields:
 1. number of different amino acids appearing in in this column of the alignment
 2. their type
- rho ET score - the smaller this value, the lesser variability of this position across the branches of the tree (and, presumably, the greater the importance for the protein)
- cvg coverage - percentage of the residues on the structure which have this rho or smaller
- gaps percentage of gaps in this column

4.2 Color schemes used

The following color scheme is used in figures with residues colored by cluster size: black is a single-residue cluster; clusters composed of more than one residue colored according to this hierarchy (ordered by descending size): red, blue, yellow, green, purple, azure, turquoise, brown, coral, magenta, LightSalmon, SkyBlue, violet, gold, bisque, LightSlateBlue, orchid, RosyBrown, MediumAquamarine, DarkOliveGreen, CornflowerBlue, grey55, burlywood, LimeGreen, tan, DarkOrange, DeepPink, maroon, BlanchedAlmond.

The colors used to distinguish the residues by the estimated evolutionary pressure they experience can be seen in Fig. 12.

4.3 Credits

4.3.1 Alistat *alistat* reads a multiple sequence alignment from the file and shows a number of simple statistics about it. These statistics include the format, the number of sequences, the total number of residues, the average and range of the sequence lengths, and the alignment length (e.g. including gap characters). Also shown are some percent identities. A percent pairwise alignment identity is defined as $(\text{idents} / \text{MIN}(\text{len1}, \text{len2}))$ where *idents* is the number of exact identities and *len1*, *len2* are the unaligned lengths of the two sequences. The “average percent identity”, “most related pair”, and “most unrelated pair” of the alignment are the average, maximum, and minimum of all $(N)(N-1)/2$ pairs, respectively. The “most distant seq” is calculated by finding the maximum pairwise identity (best relative) for all *N* sequences, then finding the minimum of these *N* numbers (hence, the most outlying sequence). *alistat* is copyrighted

by HHMI/Washington University School of Medicine, 1992-2001, and freely distributed under the GNU General Public License.

4.3.2 CE To map ligand binding sites from different source structures, `report_maker` uses the CE program: <http://cl.sdsc.edu/>. Shindyalov IN, Bourne PE (1998) "Protein structure alignment by incremental combinatorial extension (CE) of the optimal path". *Protein Engineering* 11(9) 739-747.

4.3.3 DSSP In this work a residue is considered solvent accessible if the DSSP program finds it exposed to water by at least 10\AA^2 , which is roughly the area needed for one water molecule to come in the contact with the residue. DSSP is copyrighted by W. Kabsch, C. Sander and MPI-MF, 1983, 1985, 1988, 1994 1995, CMBI version by Elmar.Krieger@cmbi.kun.nl November 18,2002,

<http://www.cmbi.kun.nl/gv/dssp/descrip.html>.

4.3.4 HSSP Whenever available, `report_maker` uses HSSP alignment as a starting point for the analysis (sequences shorter than 75% of the query are taken out, however); R. Schneider, A. de Daruvar, and C. Sander. "The HSSP database of protein structure-sequence alignments." *Nucleic Acids Res.*, 25:226-230, 1997.

<http://swift.cmbi.kun.nl/swift/hssp/>

4.3.5 LaTeX The text for this report was processed using L^AT_EX; Leslie Lamport, "LaTeX: A Document Preparation System Addison-Wesley," Reading, Mass. (1986).

4.3.6 Muscle When making alignments "from scratch", `report_maker` uses Muscle alignment program: Edgar, Robert C. (2004), "MUSCLE: multiple sequence alignment with high accuracy and high throughput." *Nucleic Acids Research* 32(5), 1792-97.

<http://www.drive5.com/muscle/>

4.3.7 Pymol The figures in this report were produced using Pymol. The scripts can be found in the attachment. Pymol is an open-source application copyrighted by DeLano Scientific LLC (2005). For more information about Pymol see <http://pymol.sourceforge.net/>. (Note for Windows users: the attached package needs to be unzipped for Pymol to read the scripts and launch the viewer.)

4.4 Note about ET Viewer

Dan Morgan from the Lichtarge lab has developed a visualization tool specifically for viewing trace results. If you are interested, please visit:

<http://mammoth.bcm.tmc.edu/traceview/>

The viewer is self-unpacking and self-installing. Input files to be used with ETV (extension .etvx) can be found in the attachment to the main report.

4.5 Citing this work

The method used to rank residues and make predictions in this report can be found in Mihalek, I., I. Reš, O. Lichtarge. (2004). "A Family of Evolution-Entropy Hybrid Methods for Ranking of Protein Residues by Importance" *J. Mol. Bio.* **336**: 1265-82. For the original version of ET see O. Lichtarge, H.Bourne and F. Cohen (1996). "An Evolutionary Trace Method Defines Binding Surfaces Common to Protein Families" *J. Mol. Bio.* **257**: 342-358.

`report_maker` itself is described in Mihalek I., I. Res and O. Lichtarge (2006). "Evolutionary Trace Report Maker: a new type of service for comparative analysis of proteins." *Bioinformatics* **22**:1656-7.

4.6 About report_maker

`report_maker` was written in 2006 by Ivana Mihalek. The 1D ranking visualization program was written by Ivica Reš. `report_maker` is copyrighted by Lichtarge Lab, Baylor College of Medicine, Houston.

4.7 Attachments

The following files should accompany this report:

- 3ecsC.complex.pdb - coordinates of 3ecsC with all of its interacting partners
- 3ecsC.etvx - ET viewer input file for 3ecsC
- 3ecsC.cluster_report.summary - Cluster report summary for 3ecsC
- 3ecsC.ranks - Ranks file in sequence order for 3ecsC
- 3ecsC.clusters - Cluster descriptions for 3ecsC
- 3ecsC.msf - the multiple sequence alignment used for the chain 3ecsC
- 3ecsC.descr - description of sequences used in 3ecsC msf
- 3ecsC.ranks_sorted - full listing of residues and their ranking for 3ecsC
- 3ecsC.3ecsE.if.pml - Pymol script for Figure 5
- 3ecsC.cbcvg - used by other 3ecsC – related pymol scripts
- 3ecsC.3ecsFSO4506.if.pml - Pymol script for Figure 6
- 3ecsC.3ecsF.if.pml - Pymol script for Figure 7
- 3ecsC.3ecsA.if.pml - Pymol script for Figure 8
- 3ecsC.3ecsCSO4503.if.pml - Pymol script for Figure 9