


3eh0

Evolutionary trace report by **report_maker**

August 24, 2009



4.3.1	Alistat	11
4.3.2	CE	11
4.3.3	DSSP	11
4.3.4	HSSP	11
4.3.5	LaTex	11
4.3.6	Muscle	11
4.3.7	Pymol	11
4.4	Note about ET Viewer	11
4.5	Citing this work	11
4.6	About report_maker	11
4.7	Attachments	11

CONTENTS

1 Introduction

2 Chain 3eh0A

- 2.1 P21645 overview
- 2.2 Multiple sequence alignment for 3eh0A
- 2.3 Residue ranking in 3eh0A
- 2.4 Top ranking residues in 3eh0A and their position on the structure
 - 2.4.1 Clustering of residues at 25% coverage.
 - 2.4.2 Overlap with known functional surfaces at 25% coverage.
 - 2.4.3 Possible novel functional surfaces at 25% coverage.

3 Notes on using trace results

- 3.1 Coverage
- 3.2 Known substitutions
- 3.3 Surface
- 3.4 Number of contacts
- 3.5 Annotation
- 3.6 Mutation suggestions

4 Appendix

- 4.1 File formats
- 4.2 Color schemes used
- 4.3 Credits

1 INTRODUCTION

From the original Protein Data Bank entry (PDB id 3eh0):

Title: Crystal structure of lpxd from escherichia coli

Compound: Mol id: 1; molecule: udp-3-o-[3-hydroxymyristoyl] glucosamine n- acyltransferase; chain: a, b, c; synonym: protein fir, rifampicin resistance protein; ec: 2.3.1.-; engineered: yes

Organism, scientific name: Escherichia Coli;

3eh0 contains a single unique chain 3eh0A (341 residues long) and its homologues 3eh0C and 3eh0B.

1 2 CHAIN 3EH0A

2.1 P21645 overview

- 1 From SwissProt, id P21645, 97% identical to 3eh0A:
- 1 **Description:** UDP-3-O-[3-hydroxymyristoyl] glucosamine N-acyltransferase (EC 2.3.1.-) (FirA protein) (Rifampicin resistance protein).
- 1 **Organism, scientific name:** Escherichia coli.
- 2 **Taxonomy:** Bacteria; Proteobacteria; Gammaproteobacteria; Enterobacteriales; Enterobacteriaceae; Escherichia.
- 2 **Catalytic activity:** UDP-3-O-(3-hydroxytetradecanoyl)glucosamine + (R)-3-hydroxytetradecanoyl-[acyl-carrier protein] = UDP-2,3-bis(3- hydroxytetradecanoyl)glucosamine + [acyl-carrier protein].
- 7 **Pathway:** Lipid A biosynthesis; third step.
- 10 **Interaction:**
- 10 **Miscellaneous:** The mutants firA200, firA201 and omsA confer temperature sensitivity; firA200 reverses the rifampicin resistance of rpoB mutants.
- 10 **Similarity:** Belongs to the transferase hexapeptide repeat family. LpxD subfamily.
- 10 **Caution:** Was originally thought to be involved in transcription.
- 10 **About:** This Swiss-Prot entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use as long as its content is in no way modified and this statement is not removed.
- 11




Fig. 1. Residues 1-170 in 3eh0A colored by their relative importance. (See Appendix, Fig.9, for the coloring scheme.)




Fig. 2. Residues 171-341 in 3eh0A colored by their relative importance. (See Appendix, Fig.9, for the coloring scheme.)

2.2 Multiple sequence alignment for 3eh0A

For the chain 3eh0A, the alignment 3eh0A.msf (attached) with 592 sequences was used. The alignment was downloaded from the HSSP database, and fragments shorter than 75% of the query as well as duplicate sequences were removed. It can be found in the attachment to this report, under the name of 3eh0A.msf. Its statistics, from the *alistat* program are the following:

```

Format:                MSF
Number of sequences:   592
Total number of residues: 195974
Smallest:              258
Largest:              341
Average length:       331.0
Alignment length:     341
Average identity:      36%
Most related pair:    99%
Most unrelated pair:  16%
Most distant seq:     33%

```

Furthermore, <1% of residues show as conserved in this alignment.

The alignment consists of <1% eukaryotic (<1% plantae), and 9% prokaryotic sequences. (Descriptions of some sequences were not readily available.) The file containing the sequence descriptions can be found in the attachment, under the name 3eh0A.descr.

2.3 Residue ranking in 3eh0A

The 3eh0A sequence is shown in Figs. 1–2, with each residue colored according to its estimated importance. The full listing of residues in 3eh0A can be found in the file called 3eh0A.ranks.sorted in the attachment.

2.4 Top ranking residues in 3eh0A and their position on the structure

In the following we consider residues ranking among top 25% of residues in the protein. Figure 3 shows residues in 3eh0A colored by their importance: bright red and yellow indicate more conserved/important residues (see Appendix for the coloring scheme). A Pymol script for producing this figure can be found in the attachment.

2.4.1 Clustering of residues at 25% coverage. Fig. 4 shows the top 25% of all residues, this time colored according to clusters they




Fig. 3. Residues in 3eh0A, colored by their relative importance. Clockwise: front, back, top and bottom views.

belong to. The clusters in Fig.4 are composed of the residues listed in Table 1.

Table 1.		
cluster color	size	member residues
red	68	149, 167, 170, 173, 175, 177, 178 179, 181, 182, 183, 184, 185, 186 192, 194, 195, 196, 197, 199, 201 204, 207, 208, 209, 210, 211, 212 215, 216, 217, 218, 219, 220, 223 225, 228, 230, 231, 232, 233, 234 235, 236, 237, 238, 239, 240, 241 244, 247, 250, 251, 252, 254, 256 257, 258, 262, 265, 268, 269, 270 272, 275, 276, 287, 291

continued in next column




Fig. 4. Residues in 3eh0A, colored according to the cluster they belong to: red, followed by blue and yellow are the largest clusters (see Appendix for the coloring scheme). Clockwise: front, back, top and bottom views. The corresponding Pymol script is attached.

Table 1. continued		
cluster color	size	member residues
blue	8	31, 34, 41, 56, 79, 80, 83, 84
yellow	2	104, 119
green	2	304, 306

Table 1. Clusters of top ranking residues in 3eh0A.

2.4.2 Overlap with known functional surfaces at 25% coverage. The name of the ligand is composed of the source PDB identifier and the heteroatom name used in that file.

Interface with 3eh0B. Table 2 lists the top 25% of residues at the interface with 3eh0B. The following table (Table 3) suggests possible disruptive replacements for these residues (see Section 3.6).

Table 2.					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
216	D	D(97)HE NC	0.03	12/0	3.02
183	F	F(97) Y(2)H	0.04	83/5	3.43
197	Q	Q(81) S(4)	0.07	2/2	4.49
181	D	H(12)IN E(12) D(82)RQ	0.08	19/9	3.56

continued in next column

Table 2. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
254	I	PHC(1) A(1)SNF G(77) I(10) T(2) V(3) A(4)NSL K	0.09	4/0	4.56
272	V	G(68) V(11) A(9) M(3) N(1) I(1) S(1) C(1)QDK TP	0.09	10/0	3.68
194	K	K(83) H(1) R(8) A(1)E T(1) P(1)G Q(1)DSM L	0.10	30/10	2.75
185	Y	F(55) Y(36)I L(3) N(1)AHC W	0.11	34/3	2.55
236	Q	Q(80) H(13) M(5)LV	0.12	44/6	3.16
175	G	G(78)K N(14) S(5)HCF AQT	0.13	2/2	3.71
235	C	I(18) V(67) C(11)AM L(1)ST	0.13	11/11	3.29
195	I	I(71) V(13) M(6) L(3) F(1)TYK GARNs	0.14	30/3	3.23
234	Q	Q(35) L(59) H(2) M(1)FIS	0.14	10/9	3.15
306	P	A(6)	0.14	18/6	3.71

continued in next column

Table 2. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
196	P	P(82)HI V(1) G(1)WQ S(1)E T(1) F(1).LN K A(6) P(66) H(4) V(3) E(6) F(1)Q Y(3)N D(1) K(2) L(2)STM I	0.16	37/5	3.69
212	C	N(67) G(8) C(15) S(4) L(1)HAF QME	0.17	10/9	3.01
177	V	V(74) Q(1) I(6) R(7) H(1) K(1) S(3) T(1) C(1)A	0.18	19/0	3.61
186	A	A(64) I(6) V(14) E(1)Q S(2) R(2) K(1) T(2)LM Y(2)H F(1)D	0.18	8/3	4.03
219	A	A(53) S(9) T(22) V(6) R(2) K(2)LN G(1)DM	0.19	4/3	3.75
304	G	G(77) Y(1)	0.20	17/17	2.83

continued in next column

Table 2. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
192	W	A(6)H I(1) S(2) T(1)L D(1) N(1) F(1) V(3)M.E PK W(49). Y(13) H(19) F(3) I(2) M(2) N(2) L(2) R(1)QVP T	0.22	54/1	3.17
220	L	L(45) M(17) V(8) I(12)G F(10) S(1) T(1) Y(1)WNA	0.23	44/1	3.73
291	V	V(72) I(20)K L(2)A G(1).RS F	0.24	9/4	3.92

Table 2. The top 25% of residues in 3eh0A at the interface with 3eh0B. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 3.		
res	type	disruptive mutations
216	D	(R)(FW)(H)(Y)
183	F	(KE)(Q)(D)(T)
197	Q	(Y)(T)(FWH)(CG)
181	D	(R)(H)(FW)(Y)
254	I	(R)(Y)(H)(E)
272	V	(Y)(R)(KEH)(D)
194	K	(Y)(FW)(T)(CHG)
185	Y	(K)(Q)(E)(R)
236	Q	(Y)(T)(H)(FW)

continued in next column

res	type	disruptive mutations
175	G	(E)(R)(K)(D)
235	C	(R)(K)(E)(H)
195	I	(YR)(H)(TE)(K)
234	Q	(Y)(T)(H)(FW)
306	P	(YR)(H)(T)(E)
196	P	(R)(Y)(H)(T)
212	C	(R)(KE)(H)(FW)
177	V	(Y)(E)(R)(K)
186	A	(Y)(R)(K)(E)
219	A	(Y)(R)(E)(K)
304	G	(R)(K)(E)(H)
192	W	(E)(K)(T)(D)
220	L	(R)(Y)(KH)(E)
291	V	(Y)(E)(R)(K)

Table 3. List of disruptive mutations for the top 25% of residues in 3eh0A, that are at the interface with 3eh0B.




Fig. 5. Residues in 3eh0A, at the interface with 3eh0B, colored by their relative importance. 3eh0B is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 3eh0A.)

Figure 5 shows residues in 3eh0A colored by their importance, at the interface with 3eh0B.

Interface with 3eh0C. Table 4 lists the top 25% of residues at the interface with 3eh0C. The following table (Table 5) suggests possible disruptive replacements for these residues (see Section 3.6).

res	type	subst's (%)	cvg	noc/ bb	dist (Å)
232	D	D(99)G	0.01	9/0	4.38
233	N	N(96) S(1) D(1)AH	0.04	38/5	3.04
269	G	G(95) A(4)PS	0.05	1/1	4.76
41	F	F(82) L(3) Y(10)C M(1)V.A	0.08	3/0	3.82
230	I	K(71) I(19) R(5)VYM CFHWQ	0.09	40/1	3.22
208	E	E(87) D(7) R(1)FVS Q(1)GTK HN	0.10	26/0	2.55
175	G	G(78)K N(14) S(5)HCF AQT	0.13	2/2	4.27
234	Q	Q(35) L(59) H(2) M(1)FIS	0.14	47/0	3.09
306	P	A(6) P(82)HI V(1) G(1)WQ S(1)E T(1) F(1).LN K	0.14	23/14	3.32
228	G	G(75)S L(1) N(10) Y(2) C(2)W D(3)F H(1)QRM	0.15	4/4	3.43
84	A	A(75) V(5) S(6) Q(1) G(2) P(1) N(1)R	0.16	7/3	4.06

continued in next column

Table 4. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
211	A	T(2)YCF MI(1)L V(1) A(78) S(12) C(5)	0.16	11/5	3.75
83	Y	T(1)GLM Y(21) F(58) L(7) M(8) W(1)AST QVIR	0.17	19/0	3.77
212	C	N(67) G(8) C(15) S(4) L(1)HAF QME	0.17	15/2	3.70
287	G	G(44) A(48) P(1) S(3)T.Y HLV	0.17	2/2	3.85
173	Q	Q(24) H(60)R L(4)F N(2) Y(2)DSG CEK.	0.19	36/2	2.70
80	Y	Y(61) F(3) K(2) R(14) H(5) S(3) D(2)N Q(3)A E(1)LVI PMT	0.20	7/0	3.52
304	G	G(77) Y(1) A(6)H I(1) S(2) T(1)L D(1) N(1) F(1) V(3)M.E PK	0.20	5/5	3.90
247	T	T(45)	0.21	3/3	4.42

continued in next column

Table 4. continued					
res	type	subst's (%)	cvg	noc/ bb	dist (Å)
270	A	C(32) S(15) N(2)AI V(2)GY G(10) K(3) Q(50) A(22) N(3) H(1)C S(2) M(1)E R(2)DL	0.21	8/3	3.36
252	G	C(17) Q(53) G(13) L(1) K(2)A N(2) S(3) H(2)VYT FM	0.22	4/4	4.24
251	G	A(39) G(41) S(19)	0.23	3/3	4.04

Table 4. The top 25% of residues in 3eh0A at the interface with 3eh0C. (Field names: res: residue number in the PDB entry; type: amino acid type; substs: substitutions seen in the alignment; with the percentage of each type in the bracket; noc/bb: number of contacts with the ligand, with the number of contacts realized through backbone atoms given in the bracket; dist: distance of closest approach to the ligand.)

Table 5.		
res	type	disruptive mutations
232	D	(R)(FWH)(K)(Y)
233	N	(Y)(R)(FWH)(T)
269	G	(R)(K)(E)(H)
41	F	(K)(E)(QD)(R)
230	I	(Y)(R)(T)(E)
208	E	(FW)(H)(Y)(R)
175	G	(E)(R)(K)(D)
234	Q	(Y)(T)(H)(FW)
306	P	(YR)(H)(T)(E)
228	G	(E)(KR)(D)(Q)
84	A	(R)(E)(Y)(K)
211	A	(R)(K)(YE)(H)
83	Y	(K)(Q)(ER)(M)
212	C	(R)(KE)(H)(FW)
287	G	(KR)(E)(Q)(HD)

continued in next column

Table 5. continued		
res	type	disruptive mutations
173	Q	(Y) (FW) (H) (T)
80	Y	(K) (R) (Q) (EM)
304	G	(R) (K) (E) (H)
247	T	(R) (K) (H) (Q)
270	A	(Y) (R) (E) (H)
252	G	(E) (R) (K) (D)
251	G	(KR) (E) (QH) (FMW)

Table 5. List of disruptive mutations for the top 25% of residues in 3eh0A, that are at the interface with 3eh0C.




Fig. 7. A possible active surface on the chain 3eh0A.




Fig. 6. Residues in 3eh0A, at the interface with 3eh0C, colored by their relative importance. 3eh0C is shown in backbone representation (See Appendix for the coloring scheme for the protein chain 3eh0A.)

Figure 6 shows residues in 3eh0A colored by their importance, at the interface with 3eh0C.

2.4.3 Possible novel functional surfaces at 25% coverage. One group of residues is conserved on the 3eh0A surface, away from (or substantially larger than) other functional sites and interfaces recognizable in PDB entry 3eh0. It is shown in Fig. 7. The residues belonging to this surface "patch" are listed in Table 6, while Table 7 suggests possible disruptive replacements for these residues (see Section 3.6).

Table 6.			
res	type	substitutions(%)	cvg
41	F	F(82)L(3)Y(10)C M(1)V.A	0.08

continued in next column

Table 6. continued			
res	type	substitutions(%)	cvg
56	A	A(93)CS(2)IPVG. QMEK	0.08
34	A	A(84)G(8)QS(2) V(1)PC.IH	0.11
79	P	P(81)A(11)S(2)Q V(2)KGNETH	0.13
84	A	A(75)V(5)S(6) Q(1)G(2)P(1) N(1)RT(2)YCFM I(1)L	0.16
83	Y	Y(21)F(58)L(7) M(8)W(1)ASTQVIR	0.17
80	Y	Y(61)F(3)K(2) R(14)H(5)S(3) D(2)NQ(3)AE(1)L VIPMT	0.20
31	M	L(54)I(22)M(9) F(2)P(4)A(2) V(2)W.YSH	0.24

Table 6. Residues forming surface "patch" in 3eh0A.

Table 7.		
res	type	disruptive mutations
41	F	(K) (E) (QD) (R)
56	A	(Y) (R) (H) (KE)
34	A	(R) (YE) (K) (H)

continued in next column

Table 7. continued		
res	type	disruptive mutations
79	P	(Y) (R) (H) (T)
84	A	(R) (E) (Y) (K)
83	Y	(K) (Q) (ER) (M)
80	Y	(K) (R) (Q) (EM)
31	M	(Y) (T) (R) (H)

Table 7. Disruptive mutations for the surface patch in 3eh0A.

Another group of surface residues is shown in Fig.8. The right panel shows (in blue) the rest of the larger cluster this surface belongs to.




Fig. 8. Another possible active surface on the chain 3eh0A. The larger cluster it belongs to is shown in blue.

The residues belonging to this surface "patch" are listed in Table 8, while Table 9 suggests possible disruptive replacements for these residues (see Section 3.6).

Table 8.			
res	type	substitutions(%)	cvg
257	G	G(99)A	0.00
182	G	G(98)A(1)P	0.01
210	G	G(99)Q	0.01
232	D	D(99)G	0.01
239	H	H(99)A	0.01
216	D	D(97)HENC	0.03
244	G	G(98)DNSE	0.03
258	S	S(90)G(6)C(1)TA LRQ	0.03
183	F	F(97)Y(2)H	0.04
184	G	G(93)S(3)Y(1)RK AN	0.04
233	N	N(96)S(1)D(1)AH	0.04
262	G	G(94)E(4)DKHSA	0.05
269	G	G(95)A(4)PS	0.05
276	H	H(87)N(4)Q(3) R(2)G(1)KADSY	0.05
149	G	G(95)QA(1).D(2) EHT	0.06

continued in next column

Table 8. continued			
res	type	substitutions(%)	cvg
217	R	R(94)N(2)K(1)CS A	0.06
240	N	N(87)G(8)D(1)HS QA	0.06
167	G	G(96)K(1)E.RDNA	0.07
197	Q	Q(81)S(4)H(12)I N	0.07
275	G	G(74)N(7)D(14) S(1)PQA	0.07
181	D	E(12)D(82)RQPH C(1)A(1)SNF	0.08
230	I	K(71)I(19)R(5)V YMCFWQ	0.09
254	I	G(77)I(10)T(2) V(3)A(4)NSLK	0.09
272	V	G(68)V(11)A(9) M(3)N(1)I(1) S(1)C(1)QDKTP	0.09
194	K	K(83)H(1)R(8) A(1)ET(1)P(1)G Q(1)DSML	0.10
208	E	E(87)D(7)R(1)FV SQ(1)GTKHN	0.10
131	G	A(3)G(87)KE(3) . (2)MD(1)SNQ	0.11
185	Y	F(55)Y(36)IL(3) N(1)AHCW	0.11
268	G	A(27)G(70)W(1)S T	0.11
236	Q	Q(80)H(13)M(5)L V	0.12
250	A	A(64)I(2)C(7) V(22)LSTP(1)MR	0.12
256	A	S(25)A(72)G(1)L	0.12
175	G	G(78)KN(14)S(5) HCFAQT	0.13
235	C	I(18)V(67)C(11) AML(1)ST	0.13
170	C	V(43)C(45)F(3) A(3)GS.TINL	0.14
195	I	I(71)V(13)M(6) L(3)F(1)TYKGARN S	0.14
234	Q	Q(35)L(59)H(2) M(1)FIS	0.14
228	G	G(75)SL(1)N(10) Y(2)C(2)WD(3)F H(1)QRM	0.15
238	A	A(62)G(37)S	0.15
196	P	A(6)P(66)H(4) V(3)E(6)F(1)Q Y(3)ND(1)K(2)	0.16

continued in next column

Table 8. continued			
res	type	substitutions(%)	cvg
211	A	L(2)STMI V(1)A(78)S(12) C(5)T(1)GLM	0.16
212	C	N(67)G(8)C(15) S(4)L(1)HAFQME	0.17
287	G	G(44)A(48)P(1) S(3)T.YHLV	0.17
177	V	V(74)Q(1)I(6) R(7)H(1)K(1) S(3)T(1)C(1)A	0.18
186	A	A(64)I(6)V(14) E(1)QS(2)R(2) K(1)T(2)LMY(2)H F(1)D	0.18
225	I	I(83)V(11)L(4)	0.18
173	Q	Q(24)H(60)RL(4) FN(2)Y(2)DSGCEK .	0.19
219	A	A(53)S(9)T(22) V(6)R(2)K(2)LN G(1)DM	0.19
204	G	G(54)Q(8)E(30) A(1)D(2)MHSK(1) C	0.20
209	I	I(84)V(12)L(2)M	0.21
270	A	G(10)K(3)Q(50) A(22)N(3)H(1)C S(2)M(1)ER(2)DL	0.21
192	W	W(49).Y(13) H(19)F(3)I(2) M(2)N(2)L(2) R(1)QVPT	0.22
252	G	C(17)Q(53)G(13) L(1)K(2)AN(2) S(3)H(2)VYTFM	0.22
220	L	L(45)M(17)V(8) I(12)GF(10)S(1) T(1)Y(1)WNA	0.23
231	I	L(28)I(66)V(2) M(1)F(1)T	0.23
251	G	A(39)G(41)S(19)	0.23
291	V	V(72)I(20)KL(2) AG(1).RSF	0.24
279	I	I(75)L(10)V(11) .M(1)H	0.25

Table 8. Residues forming surface "patch" in 3eh0A.

Table 9.		
res	type	disruptive mutations
257	G	(KER)(QHD)(FYMW)(N)
182	G	(R)(KE)(H)(Y)
continued in next column		

Table 9. continued		
res	type	disruptive mutations
210	G	(FEWHR)(KYD)(M)(QLPI)
232	D	(R)(FWH)(K)(Y)
239	H	(E)(QD)(TKM)(N)
216	D	(R)(FW)(H)(Y)
244	G	(R)(K)(FWH)(YM)
258	S	(R)(K)(H)(FW)
183	F	(KE)(Q)(D)(T)
184	G	(E)(KR)(FW)(D)
233	N	(Y)(R)(FWH)(T)
262	G	(R)(K)(FEW)(H)
269	G	(R)(K)(E)(H)
276	H	(E)(D)(TM)(Q)
149	G	(R)(K)(FEWH)(QM)
217	R	(YD)(E)(T)(FW)
240	N	(Y)(FWH)(R)(T)
167	G	(FWR)(H)(E)(KY)
197	Q	(Y)(T)(FWH)(CG)
275	G	(R)(H)(K)(E)
181	D	(R)(H)(FW)(Y)
230	I	(Y)(R)(T)(E)
254	I	(R)(Y)(H)(E)
272	V	(Y)(R)(KEH)(D)
194	K	(Y)(FW)(T)(CHG)
208	E	(FW)(H)(Y)(R)
131	G	(R)(H)(FW)(Y)
185	Y	(K)(Q)(E)(R)
268	G	(K)(R)(E)(Q)
236	Q	(Y)(T)(H)(FW)
250	A	(R)(Y)(KE)(H)
256	A	(R)(K)(YE)(H)
175	G	(E)(R)(K)(D)
235	C	(R)(K)(E)(H)
170	C	(R)(K)(E)(H)
195	I	(YR)(H)(TE)(K)
234	Q	(Y)(T)(H)(FW)
228	G	(E)(KR)(D)(Q)
238	A	(KR)(E)(Y)(QH)
196	P	(R)(Y)(H)(T)
211	A	(R)(K)(YE)(H)
212	C	(R)(KE)(H)(FW)
287	G	(KR)(E)(Q)(HD)
177	V	(Y)(E)(R)(K)
186	A	(Y)(R)(K)(E)
225	I	(YR)(H)(T)(KE)
173	Q	(Y)(FW)(H)(T)
219	A	(Y)(R)(E)(K)
204	G	(R)(K)(FWH)(E)
209	I	(Y)(R)(H)(T)
270	A	(Y)(R)(E)(H)
192	W	(E)(K)(T)(D)
252	G	(E)(R)(K)(D)
220	L	(R)(Y)(KH)(E)
continued in next column		

Table 9. continued		
res	type	disruptive mutations
231	I	(R) (Y) (H) (TK)
251	G	(KR) (E) (QH) (FMW)
291	V	(Y) (E) (R) (K)
279	I	(Y) (R) (T) (H)

Table 9. Disruptive mutations for the surface patch in 3eh0A.

3 NOTES ON USING TRACE RESULTS

3.1 Coverage

Trace results are commonly expressed in terms of coverage: the residue is important if its “coverage” is small - that is if it belongs to some small top percentage of residues [100% is all of the residues in a chain], according to trace. The ET results are presented in the form of a table, usually limited to top 25% percent of residues (or to some nearby percentage), sorted by the strength of the presumed evolutionary pressure. (I.e., the smaller the coverage, the stronger the pressure on the residue.) Starting from the top of that list, mutating a couple of residues should affect the protein somehow, with the exact effects to be determined experimentally.

3.2 Known substitutions

One of the table columns is “substitutions” - other amino acid types seen at the same position in the alignment. These amino acid types may be interchangeable at that position in the protein, so if one wants to affect the protein by a point mutation, they should be avoided. For example if the substitutions are “RVK” and the original protein has an R at that position, it is advisable to try anything, but RVK. Conversely, when looking for substitutions which will *not* affect the protein, one may try replacing, R with K, or (perhaps more surprisingly), with V. The percentage of times the substitution appears in the alignment is given in the immediately following bracket. No percentage is given in the cases when it is smaller than 1%. This is meant to be a rough guide - due to rounding errors these percentages often do not add up to 100%.

3.3 Surface

To detect candidates for novel functional interfaces, first we look for residues that are solvent accessible (according to DSSP program) by at least 10\AA^2 , which is roughly the area needed for one water molecule to come in the contact with the residue. Furthermore, we require that these residues form a “cluster” of residues which have neighbor within 5\AA from any of their heavy atoms.

Note, however, that, if our picture of protein evolution is correct, the neighboring residues which *are not* surface accessible might be equally important in maintaining the interaction specificity - they should not be automatically dropped from consideration when choosing the set for mutagenesis. (Especially if they form a cluster with the surface residues.)

3.4 Number of contacts

Another column worth noting is denoted “noc/bb”; it tells the number of contacts heavy atoms of the residue in question make across the interface, as well as how many of them are realized through the backbone atoms (if all or most contacts are through the backbone,

mutation presumably won’t have strong impact). Two heavy atoms are considered to be “in contact” if their centers are closer than 5\AA .

3.5 Annotation

If the residue annotation is available (either from the pdb file or from other sources), another column, with the header “annotation” appears. Annotations carried over from PDB are the following: site (indicating existence of related site record in PDB), S-S (disulfide bond forming residue), hb (hydrogen bond forming residue, jb (james bond forming residue), and sb (for salt bridge forming residue).

3.6 Mutation suggestions

Mutation suggestions are completely heuristic and based on complementarity with the substitutions found in the alignment. Note that they are meant to be **disruptive** to the interaction of the protein with its ligand. The attempt is made to complement the following properties: small [*AVGSTC*], medium [*LPNQDEMIK*], large [*WFYHR*], hydrophobic [*LPVAMWFI*], polar [*GTCY*]; positively [*KHR*], or negatively [*DE*] charged, aromatic [*WFYH*], long aliphatic chain [*EKRQM*], OH-group possession [*SDETY*], and NH₂ group possession [*NQRK*]. The suggestions are listed according to how different they appear to be from the original amino acid, and they are grouped in round brackets if they appear equally disruptive. From left to right, each bracketed group of amino acid types resembles more strongly the original (i.e. is, presumably, less disruptive) These suggestions are tentative - they might prove disruptive to the fold rather than to the interaction. Many researcher will choose, however, the straightforward alanine mutations, especially in the beginning stages of their investigation.

4 APPENDIX

4.1 File formats

Files with extension “ranks_sorted” are the actual trace results. The fields in the table in this file:

- `alignment#` number of the position in the alignment
- `residue#` residue number in the PDB file
- `type` amino acid type
- `rank` rank of the position according to older version of ET
- `variability` has two subfields:
 1. number of different amino acids appearing in in this column of the alignment
 2. their type
- `rho` ET score - the smaller this value, the lesser variability of this position across the branches of the tree (and, presumably, the greater the importance for the protein)
- `cvg` coverage - percentage of the residues on the structure which have this rho or smaller
- `gaps` percentage of gaps in this column

4.2 Color schemes used

The following color scheme is used in figures with residues colored by cluster size: black is a single-residue cluster; clusters composed of more than one residue colored according to this hierarchy (ordered by descending size): red, blue, yellow, green, purple, azure, turquoise, brown, coral, magenta, LightSalmon, SkyBlue, violet, gold, bisque, LightSlateBlue, orchid, RosyBrown, MediumAquamarine,




Fig. 9. Coloring scheme used to color residues by their relative importance.

DarkOliveGreen, CornflowerBlue, grey55, burlywood, LimeGreen, tan, DarkOrange, DeepPink, maroon, BlanchedAlmond.

The colors used to distinguish the residues by the estimated evolutionary pressure they experience can be seen in Fig. 9.

4.3 Credits

4.3.1 Alistat *alistat* reads a multiple sequence alignment from the file and shows a number of simple statistics about it. These statistics include the format, the number of sequences, the total number of residues, the average and range of the sequence lengths, and the alignment length (e.g. including gap characters). Also shown are some percent identities. A percent pairwise alignment identity is defined as $(\text{idents} / \text{MIN}(\text{len1}, \text{len2}))$ where *idents* is the number of exact identities and *len1*, *len2* are the unaligned lengths of the two sequences. The "average percent identity", "most related pair", and "most unrelated pair" of the alignment are the average, maximum, and minimum of all $(N)(N-1)/2$ pairs, respectively. The "most distant seq" is calculated by finding the maximum pairwise identity (best relative) for all *N* sequences, then finding the minimum of these *N* numbers (hence, the most outlying sequence). *alistat* is copyrighted by HHMI/Washington University School of Medicine, 1992-2001, and freely distributed under the GNU General Public License.

4.3.2 CE To map ligand binding sites from different source structures, *report_maker* uses the CE program: <http://cl.sdsc.edu/>. Shindyalov IN, Bourne PE (1998) "Protein structure alignment by incremental combinatorial extension (CE) of the optimal path". *Protein Engineering* 11(9) 739-747.

4.3.3 DSSP In this work a residue is considered solvent accessible if the DSSP program finds it exposed to water by at least 10\AA^2 , which is roughly the area needed for one water molecule to come in the contact with the residue. DSSP is copyrighted by W. Kabsch, C. Sander and MPI-MF, 1983, 1985, 1988, 1994 1995, CMBI version by Elmar.Krieger@cmbi.kun.nl November 18,2002,

<http://www.cmbi.kun.nl/gv/dssp/descrip.html>.

4.3.4 HSSP Whenever available, *report_maker* uses HSSP alignment as a starting point for the analysis (sequences shorter than 75% of the query are taken out, however); R. Schneider, A. de

Daruvar, and C. Sander. "The HSSP database of protein structure-sequence alignments." *Nucleic Acids Res.*, 25:226-230, 1997.

<http://swift.cmbi.kun.nl/swift/hssp/>

4.3.5 LaTeX The text for this report was processed using \LaTeX ; Leslie Lamport, "LaTeX: A Document Preparation System Addison-Wesley," Reading, Mass. (1986).

4.3.6 Muscle When making alignments "from scratch", *report_maker* uses Muscle alignment program: Edgar, Robert C. (2004), "MUSCLE: multiple sequence alignment with high accuracy and high throughput." *Nucleic Acids Research* 32(5), 1792-97.

<http://www.drive5.com/muscle/>

4.3.7 Pymol The figures in this report were produced using Pymol. The scripts can be found in the attachment. Pymol is an open-source application copyrighted by DeLano Scientific LLC (2005). For more information about Pymol see <http://pymol.sourceforge.net/>. (Note for Windows users: the attached package needs to be unzipped for Pymol to read the scripts and launch the viewer.)

4.4 Note about ET Viewer

Dan Morgan from the Lichtarge lab has developed a visualization tool specifically for viewing trace results. If you are interested, please visit:

<http://mammoth.bcm.tmc.edu/traceview/>

The viewer is self-unpacking and self-installing. Input files to be used with ETV (extension .etvx) can be found in the attachment to the main report.

4.5 Citing this work

The method used to rank residues and make predictions in this report can be found in Mihalek, I., I. Reš, O. Lichtarge. (2004). "A Family of Evolution-Entropy Hybrid Methods for Ranking of Protein Residues by Importance" *J. Mol. Bio.* **336**: 1265-82. For the original version of ET see O. Lichtarge, H.Bourne and F. Cohen (1996). "An Evolutionary Trace Method Defines Binding Surfaces Common to Protein Families" *J. Mol. Bio.* **257**: 342-358.

report_maker itself is described in Mihalek I., I. Res and O. Lichtarge (2006). "Evolutionary Trace Report Maker: a new type of service for comparative analysis of proteins." *Bioinformatics* **22**:1656-7.

4.6 About report_maker

report_maker was written in 2006 by Ivana Mihalek. The 1D ranking visualization program was written by Ivica Reš. **report_maker** is copyrighted by Lichtarge Lab, Baylor College of Medicine, Houston.

4.7 Attachments

The following files should accompany this report:

- 3eh0A.complex.pdb - coordinates of 3eh0A with all of its interacting partners
- 3eh0A.etvx - ET viewer input file for 3eh0A
- 3eh0A.cluster_report.summary - Cluster report summary for 3eh0A

- 3eh0A.ranks - Ranks file in sequence order for 3eh0A
- 3eh0A.clusters - Cluster descriptions for 3eh0A
- 3eh0A.msf - the multiple sequence alignment used for the chain 3eh0A
- 3eh0A.descr - description of sequences used in 3eh0A msf
- 3eh0A.ranks_sorted - full listing of residues and their ranking for 3eh0A
- 3eh0A.3eh0B.if.pml - Pymol script for Figure 5
- 3eh0A.cbcvg - used by other 3eh0A – related pymol scripts
- 3eh0A.3eh0C.if.pml - Pymol script for Figure 6